

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/322931527>

# Ecosystem carbon sequestration of different land-uses of the lowland wet zone: A case study from Waga area, Kalutara District, Sri Lanka

Conference Paper · February 2018

CITATIONS

0

READS

27

3 authors, including:



**Janaka Kuruppuarachchi**

The Open University of Sri Lanka

93 PUBLICATIONS 78 CITATIONS

[SEE PROFILE](#)



**Gamini Senevirate**

National Institute of Fundamental Studies - Sri Lanka

129 PUBLICATIONS 1,182 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Urban flora in Colombo City ,Sri Lanka [View project](#)



BOOK: Agro-Environmental Sustainability (Two volumes from Springer-2017) [View project](#)

IRSUWU 2018



විද්‍යා, තාක්ෂණ හා පර්යේෂණ අමාත්‍යාංශය  
விஞ்ஞான, தொழில்நுட்பவியல் மற்றும் ஆராய்ச்சி அமைச்சு  
Ministry of Science, Technology and Research



# 2<sup>nd</sup> International Research Symposium

"Emerging Technologies for an Innovative Green Economy"

February 1–2, 2018

Uva Wellassa University  
Badulla  
Sri Lanka



**IRSUWU 2018**

## **2<sup>nd</sup> International Research Symposium**

**“Emerging Technologies for an Innovative Green Economy”**

**February 1 – 2, 2018**

**Uva Wellassa University  
Badulla  
Sri Lanka**

## Ecosystem Carbon Sequestration of Different Land-uses of the Lowland Wet Zone: A Case Study from Waga Area, Kalutara District, Sri Lanka

K.A.J.M. Kuruppuarachchi<sup>1\*</sup>, B.D. Madurapperuma<sup>2</sup> and G. Seneviratne<sup>3</sup>

<sup>1</sup>Department of Botany, The Open University of Sri Lanka, Nawala, Nugegoda, Sri Lanka

<sup>2</sup>Department of Forestry and Wildland Resources, Humboldt State University, Arcata, California, USA

<sup>3</sup>National Institute of Fundamental Studies (NIFS), Hantana Road, Kandy, Sri Lanka

This study examines plant above-ground carbon and soil carbon stocks of different land-use types with the same climate and geology in Waga area i.e., natural forests (NF), home gardens (HG), coconut plantations (CP), pine and Araucaria mixed plantation forest (PAP), rubber plantation (RP), rambutan plantation (RAP), pineapple plantation (PP) and tea plantation (TP). Five representative sampling sites viz., 20 m × 20 m were selected in each land-use for floristic survey and to make composite soil core sampling (20 samples mixed together) up to a 30 cm depth. Plant parameters, such as DBH, height and physicochemical properties of soil samples were evaluated. The normalized difference vegetation index (NDVI) was calculated using 2017 Landsat 8 image at 30 m spatial resolution, which was acquired during leaf-on season (i.e. mid January to mid March), to recognize the vegetation health. The study revealed ecosystem carbon and soil C densities in the order NF (625 and 52 t C ha<sup>-1</sup>), PAP (290 and 38 t C ha<sup>-1</sup>), HG (199 and 21 t C ha<sup>-1</sup>), RP (188 and 17 t C ha<sup>-1</sup>), CP (167 and 25 t C ha<sup>-1</sup>), RAP (167 and 28 t C ha<sup>-1</sup>), PP (40 and 38 t C ha<sup>-1</sup>) and TP (15 and 15 t C ha<sup>-1</sup>), respectively. NDVI values of the land-uses were RP (0.623), HG (0.618), NF (0.615), CP (0.611), PP (0.592), TP (0.576), RAP (0.562), and PAP (0.556). The highest soil moisture accumulation was reported in NF (17.7%) and RP (16.38%). Soil pH values were mainly acidic (< 5) for all land-uses. High total soil N concentrations were observed in both NF (0.18%) and PAP (0.14%). Higher amounts of soil nitrate were recorded in RAP (15.55 µg g<sup>-1</sup> soil), NF (11.45 µg g<sup>-1</sup> soil) and PP (9.32 µg g<sup>-1</sup> soil). Soil total P concentrations were relatively low in NF (0.041%), PAP (0.042%) and RP (0.045%), and it is an obvious fact for tropical land-uses with perennial vegetation. This study revealed that the land-use types, such as NF, PAP, HG and RP were good carbon reserves in tropical lowland wet zone of Sri Lanka. Further investigations on micro carbon cycles of each land-use are recommended for better understanding of ecosystem carbon footprint.

**Keywords:** Ecosystem carbon, Land-use types, Low land wet zone, Waga