

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/337050120>

The potential of genetic modification in enhancing cellulase production by fungal species

Conference Paper · November 2019

CITATIONS

0

READS

5

3 authors:



[Varuna Jayasinghe](#)

Horizon Campus

5 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



[Sandhya Jayasekara](#)

National Institute of Fundamental Studies - Sri Lanka

21 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



[Renuka Ratnayake](#)

National Institute of Fundamental Studies - Sri Lanka

50 PUBLICATIONS 82 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Applications of Microbes [View project](#)



Bio energy and Soil Ecosystems [View project](#)

PP-4

The potential of genetic modification in enhancing cellulase production by fungal species

Varuna Jayasinghe¹, Sandhya Jayasekara¹, Renuka Ratnayake¹

¹Bioenergy and Soil Ecosystems Research Group, National Institute of Fundamental Studies, Hantana Road, Kandy, Sri Lanka

*varuna2560@gmail.com

Shortage of fossil fuels and their detrimental environmental impacts have resulted in a need for a renewable source of energy. Celluloses, the major components of plant biomass, can be converted into sugars and fermented to produce bioethanol, which has shown potential as a sustainable alternative. Many micro-organisms, particularly fungi, have the ability to convert cellulose into sugars using cellulase enzymes. However, they lack the efficiency to produce cellulases in amounts necessary for economically feasible bioethanol production. Thus, genetic modification has been explored as a means to solve this problem. This study reviews genetic modification techniques that have successfully enhanced fungal cellulase production in comparison with isolated wild type fungi. The reviewed studies used a range of techniques to modify the genetic structure of selected fungal species, and measured total cellulase activity (FPase) by Filter Paper Assay. In one study, *Aspergillus niger* strains were exposed to different doses of γ -rays of Co⁶⁰, with survivors exposed to sequential UV irradiation. Another study incorporated *Agrobacterium*-mediated transformation to modify *Trichoderma reesei* with an amplified beta-glucosidase I gene of *Penicillium decumbens*. Constitutional overexpression of *clrB* gene in *Penicillium oxalicum* was also done, coupled with the deletion of *creA*, a gene which represses the expression of cellulase genes. FPase of the modified *Aspergillus*, *Trichoderma* and *Penicillium* strains were recorded as 11.05 U/mL, 9.87 U/mL and 6.94 U/mL respectively, while the isolated species from the present study showed FPases of 0.464 U/mL, 0.574 U/mL and 0.438 U/mL respectively. The modified strains have thus shown up to a 20-fold increase in cellulase production compared to their respective wild types. This shows the potential of genetic modification in creating efficient fungal strains for feasible large-scale bioethanol production. The application of such techniques would be valuable for the present study as well.

Keywords: *Aspergillus*, Cellulase, Fungi, Genetic modification, *Penicillium*, *Trichoderma*