



9TH ANRAP INTERNATIONAL SEMINAR
"FROM PLANT TO PATIENT: IDENTIFICATION OF
NEW DRUGS FOR METABOLIC DISORDERS (ANRAP-9)"

January 25 - 27, 2019

Jointly organized by

Dr. Panjwani Center for Molecular Medicine and Drug Research
(International Center for Chemical and Biological Sciences)

and

Asian Network of Research on Anti-diabetic Plants (ANRAP)

ANRAP BOARD

CHAIRMAN

Prof. Dr. Mohammed Mosihuzzaman
(Bangladesh)

GENERAL SECRETARY

Prof. Dr. Begum Rokeya
(Bangladesh)

MEMBERS

Prof. Dr. M. Iqbal Choudhary (Pakistan)
Prof. Dr. Khozirah Shaari (Malaysia)
Prof. Dr. Banasri Hazra (India)

ORGANIZING COMMITTEE (ANRAP-9)

PRESIDENT

Prof. Dr. Atta-ur-Rahman, FRS
Nishan-e-Imtiaz, Hilal-e-Imtiaz, Sitara-e-Imtiaz, Tamgha-e-Imtiaz
Patron-in-Chief, ICCBS

CHAIRMAN

Prof. Dr. M. Iqbal Choudhary
Hilal-e-Imtiaz, Sitara-e-Imtiaz, Tamgha-e-Imtiaz
Director, ICCBS

ORGANIZING SECRETARY

Dr. Asmat Salim
Associate Professor, ICCBS

SEMINAR COORDINATOR

Dr. Md. Hafizur Rahman
Associate Professor, ICCBS

MEMBERS

Prof. Dr. Viqar U. Ahmed, H.I., S.I.
Prof. Dr. Bina S. Siddiqui S.I., T.I.
Prof. Dr. Fatima Basha, S.I.
Prof. Dr. Khalid M. Khan, T.I., S.I.
Prof. Dr. Shaheen Faizl, S.I.
Prof. Dr. Sabira Begum
Prof. Dr. Muhammad Shaiq Ali, T.I.
Prof. Dr. Darakhshan J. Haleem
Prof. Dr. M. Iqbal Bhanger, P.P.
Prof. Dr. Farzana Shaheen
Prof. Dr. Muhammad Raza Shah, T.I.
Prof. Dr. Syed Abid Ali
Prof. Dr. Syed Ghulam Musharraf
Dr. Shabana U. Simjee
Dr. Zaheer-ul-haq Qasmi
Dr. Shakil Ahmad
Dr. Sammer Yousuf

Dr. Atia-tul-Wahab
Dr. Maqsood Chotani
Dr. Muhammad Imran Malik
Dr. Ishtiaq Ahmed Khan
Dr. Reaz Uddin
Dr. Hina Siddiqui
Dr. Ayaz Ahmad
Dr. Mudassar Azhar
Dr. Abdul Hameed
Dr. Syed Tarique Moin
Dr. Humera Jahan
Dr. Almas Jabeen
Dr. Sobia Rana
Dr. Sadia Aijaz
Dr. Irfan Khan
Dr. Naghma Hashmi
Dr. Munazza Raza

Dr. Asma Rauf
Dr. Ghulam Mustafa
Dr. Mujeeb-ur-Rahman
Dr. Seima Rasheed
Dr. Arslan Ali
Dr. Sumaira Ashraf
Dr. Maria A. Khan
Dr. Farooq Khan
Dr. Ghayas Siddiqi
Dr. Humaira Zafar
Dr. Sumaira Javed
Dr. Muhammad Rashid
Dr. Ambrina Akbar
Dr. Amna Jabbar Siddiqui
Dr. Uzma Salar
Dr. Naghma Naz
Dr. Ali Raza Shah

Dr. Zarrin Basharat
Dr. Salman Khan
Dr. Syeda Rehana Zia
Dr. M. Shoaib Malik
Dr. Muhammad Shakeel
Dr. Muhammad Irfan
Dr. Zehra Batool
Dr. Muniza Shaikh
Dr. Yan Wang

Management

Mr. Raheel Baig
Mr. Javed Riaz
Mr. Mahboob Alam

α -Amylase Enzyme inhibitory and Antioxidant Activities of Fruits of Four Sri Lankan *Artocarpus* Species

M. M. Qader^a, N. R. Amarasinghe^b, and L. Jayasinghe^a

^aNational Institute of Fundamental Studies, Kandy, Sri Lanka;

^bDepartment of Pharmacy, Faculty of Allied Health Sciences,

University of Peradeniya, Sri Lanka;

Email: nilupa16@gmail.com

Genus *Artocarpus* of the family Moraceae consists of approximately 50 species of plants distributed in Southeast Asian and Pacific regions. Some *Artocarpus* species have been extensively studied and reported to produce many secondary metabolites with diverse structures. Most of the secondary metabolites are phenolic compounds including flavonoids, stilbenoids, arilbenzofurans, neolignans etc.

In a continuation of our research work on bioactive secondary metabolites from Sri Lankan plants, present study is aimed at investigating *in vitro* α -amylase enzyme inhibitory and antioxidant activities of edible fruits of four *Artocarpus* species found in Sri Lanka. α -Amylase is a clinically important enzyme in controlling type II diabetes. Inhibition of this enzyme significantly decreases the postprandial increase of blood glucose level after a meal by delaying the carbohydrate hydrolysis and absorption. Antioxidants are capable of stabilizing or deactivating the damage caused by free radicals.

Artocarpus nobilis is the only endemic species of the genus *Artocarpus* growing in Sri Lanka. Both seeds and young fruits are edible. *A. altilis* and *A. heterophyllus* are popular staple crops and known as jack fruit and bread fruit respectively. Fruits of *A. camansi* is known as bread nut and commonly harvested from the wild as a source of food.

Dried powdered whole fruit of *A. altilis*, *A. camansi*, *A. heterophyllus* and *A. nobilis* were defatted with *n*-hexane and extracted with dichloromethane (DCM), ethyl acetate (EtOAc) and methanol (MeOH) at room temperature using a sonicator. Solvents were evaporated under reduced pressure to obtain crude extracts. Each crude extract was subjected to α -amylase enzyme inhibitory assay and DPPH radical scavenging assay using standard protocols. Both DCM and MeOH extracts of *A. nobilis* showed strong DPPH radical scavenging activity with IC₅₀ of 26 and 19 ppm respectively (positive control BHA- 19 ppm). Three extracts *A. camansi* exhibited strong antioxidant activity having IC₅₀ of 18 ppm (DCM), 12 ppm (EtOAc) and 18 ppm (MeOH). MeOH extract of *A. nobilis* and EtOAc extract of *A. camansi* showed 100% inhibition of α -amylase enzyme activity at 167 ppm (positive control acarbose 100% inhibition at 167 ppm). Results indicated that fruits of *A. altilis* and *A. camansi* contain secondary metabolites with strong DPPH radical scavenging and α -amylase inhibitory activities.