

ICSBE24\_311

## COMMUNITY APPROACH TOWARDS MODERN WATER TREATMENT TECHNOLOGY; STUDY IN ASAMODHAGAMYAYA

**J. Ketharani<sup>1\*</sup>, D.R. Samarajeewa<sup>2</sup>, R.M.L.D. Rathnayake<sup>1</sup>, S.K. Weragoda<sup>2</sup>,  
K.B.S.N. Jinadasa<sup>1,3</sup>, M.I.M. Mowjood<sup>4</sup>, J.M.P.N. Anuradha<sup>4</sup>, R. Weerasooriya<sup>5</sup>,  
S.L. Schensul<sup>6</sup>**

<sup>1</sup>Department of Civil Engineering, Faculty of Engineering, University of Peradeniya, Prof E.O.E  
Pereira Mawatha, Meewathura, Peradeniya 20400, Sri Lanka

<sup>2</sup>China Sri Lanka joint research demonstration center for water technology, Ministry of Water  
Supply, Prof E.O.E Pereira Mawatha, Meewathura, Peradeniya 20400, Sri Lanka

<sup>3</sup>Central Queensland University, Bundaberg, Queensland, Australia

<sup>4</sup>Post Graduate Institute of Agriculture (PGIA), University of Peradeniya, Old Galaha Road,  
Peradeniya 20400, Sri Lanka

<sup>5</sup>National Center for Water Quality Research, National Institute of Fundamental Studies, Kandy  
20000, Sri Lanka

<sup>6</sup>Department of Public Health Sciences, School of Medicine, University of Connecticut,  
Farmington, CT, USA

\*Correspondence E-mail: ketha94@pdn.ac.lk, TP: +94775144200

**Abstract:** Groundwater is the main source of drinking water for the majority of the people in Sri Lanka. The prevalence of Chronic Kidney Disease of Unknown etiology (CKDu) and its hypothesized link to groundwater consumption have set a barrier to its use. This has led to significant changes in the water consumption behaviour of the society. The recent identification of CKDu patients in Central provinces highlighted the need to install water treatment units in the endemic area. Consequently, five nanofiltration (NF) plants were installed within the Minipe area to provide safe water. Among these, Asomodhagamya Grama Niladhari Division (GND) was selected to study the community engagement in adapting to treated water, particularly NF-treated water, and its impact on water consumption behaviour. It was identified that the awareness of CKDu is low within the community, and consumption of NF-treated water is less compared to the other areas with high CKDu prevalence, notably North Central Province. However, the identification of CKDu patients in the community has significantly influenced water consumption behavior with more people now willing to check the quality of their groundwater. Many prefer to continue using groundwater that meets drinking water standards over NF-treated water or water treated by other systems. The reduced number of NF system beneficiaries is attributed to community structure, internal disagreements, and misconceptions about the quality of NF-treated water. Educating and advising the villagers about the risks of CKDu and the importance of safe water consumption and the proven quality of the NF-treated water will likely increase the number of beneficiaries using NF system-treated water.

**Keywords:** CKDu; Groundwater; Water Treatment; Drinking Water Quality; Nanofiltration; Community Engagement