

**POSTGRADUATE INSTITUTE OF SCIENCE
UNIVERSITY OF PERADENIYA
SRI LANKA**



**PGIS RESEARCH CONGRESS 2021
PROCEEDINGS
29th – 31st October 2021**

Copyright © 2021 by Postgraduate Institute of Science

All rights reserved. No part of this publication may be reproduced, distributed, stored in a retrieval system, and transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the publisher.

ISBN 978-955-8787-09-0

Published by

Postgraduate Institute of Science (PGIS)

University of Peradeniya

Peradeniya 20400

SRI LANKA

Printed by

Sanduni Offset Printers (Pvt) Ltd,

1/4, Sarasavi Uyana Goodshed Road,

Sarasavi Uyana, Peradeniya 20400,

SRI LANKA

Printed in the Democratic Socialist Republic of Sri Lanka

**ISOLATION AND CHARACTERIZATION OF PHENOLIC COMPOUNDS FROM
ANTIDIABETIC PLANT, *Costus speciosus***

**C.B. Gunawardhana¹, N.S. Kumar¹, N.K.B. Adikaram¹, L. Jayasinghe^{1*}, H. Araya² and
Y. Fujimoto^{1,2}**

¹National Institute of Fundamental Studies, Kandy, Sri Lanka

²School of Agriculture, Meiji University, Kawasaki, Japan

*lalith.ja@nifs.ac.lk

The medicinal plant *Costus speciosus* (Koen) Sm. or crepe ginger, *Thebu* in Sinhala, belongs to the Family *Costaceae* and inherits a range of pharmacological activities, antidiabetic effect being reported as of great importance. The plant is also well known for its antioxidant, antibacterial, anthelmintic, anticancer, anxiolytic, anti-inflammatory, antipyretic, hepato-protective and diuretic properties. Phenolic compounds are highly significant plant secondary metabolites that exert various health benefits in humans, including antioxidant effects, chemopreventive properties and anti-inflammatory activity. Therefore, the current study focused on isolating phenolic compounds in the leaf extracts of *C. speciosus*. Dried, powdered leaves were subjected to sequential extraction with hexane, ethyl acetate and methanol through ultrasound sonication. Ethyl acetate and methanol extracts were combined, re-dissolved in dichloromethane and partitioned with aqueous sodium bicarbonate followed by aqueous sodium hydroxide. Aqueous layers were re-extracted with ethyl acetate after neutralization, and the ethyl acetate extract was fractionated via silica gel column chromatography and size exclusion chromatography (Sephadex LH-20) and PTLC. Isolated compounds were characterized via ¹H NMR and ¹³C NMR spectroscopy. This approach furnished *trans*-cinnamaldehyde, *p*-coumaric acid, 4-hydroxybenzoic acid, vanillic acid, ferulic acid, indole-3-carboxaldehyde, kaempferol-3-*O*- β -D-glucopyranoside, sitosterol glucoside and (*E*)-1,3-bis(4-hydroxy-3-methoxyphenyl) prop-2-en-1-one. These compounds possess several bioactivities, including antioxidant, antibacterial and anti-inflammatory activities. Dissolving phenolic compounds into a basic aqueous layer and re-extraction with organic solvent after neutralization seems to be an efficient method to isolate phenolic compounds from plant extracts.

Financial assistance from the National Research Council, Sri Lanka (Grant No. NRC 14-13) is acknowledged.

Keywords: *Costus speciosus*, Phenolic compounds, Leaf extracts