

First record of cyanobacteria species: Cephalothrix komarekiana, from tropical Asia

Md. Fuad Hossain^{1,2,3}, R.W.T.M.R.T.K. Bowange¹, K.L.W. Kumara², D.N. Magana-Arachchi¹,

R.R. Ratnayake¹

¹National Institute of Fundamental Studies, Hantana Road, Kandy, Sri Lanka ²Department of Agricultural Biology, Faculty of Agriculture, University of Ruhuna, Sri Lanka ³Department of Biochemistry and Molecular Biology, Cono Bishwabidyalay, Savar, Dhaka, Bangladesh

ABSTRACT

Cyanobacteria play an important ecological role in every ecosystem with hidden economic and social impact. Among tropical countries, Sri Lanka holds a rich biodiversity with remarkable cyanobacterial diversity. The major objective of this study was to investigate the diversity of culturable cyanobacteria from freshwater bodies of Sri Lanka, with a phylogenetic perspective. Molecular characterization followed by phylogenetic evaluations allowed identification of new species with value added potential and revealed deeper taxonomic resolutions. A total of 74 monocultures were obtained. Rare isolates; Alkalinema pantanalense (U4), Geitlerinema sp. (U36) and Westiellopsis prolifica (U58) were also among them. Novel species Cephalothrix komarekiana was identified for the first time in Sri Lanka. This species was first reported from South American continent (Brazil and Amazonian rivers in 2015 and 2017 respectively) and Korea in 2018. The present study is next in line reporting the species first time from the tropical Asia. Higher value added potential and easy culturing of C. komarekiana suggest its economical applicability in many industries. Overall, this study provides insight on diversity, ecological distribution and industrial applicability of culturable cyanobacteria from the tropics with special reference to C. komarekiana reported from Sri Lanka.

Keywords: Cephalothrix komarekiana, Cyanobacteria, Diversity, Freshwater

1. Introduction

Cyanobacteria is an important group of microorganisms to be studied due to their nutritional simplicity, ubiquitous distribution and in some species, capacity of nitrogen fixation and toxin production.

Higher variation in morphological features among cyanobacteria even within the same species highlights the essential necessity of the contribution of morphological features in identification [1]. Branching patterns, formation of special structures such as heterocysts, akinetes, hormogonia and patterns of motility are the mostly considering properties of interest [1]. Earlier cyanobacteria identification and classification were therefore mainly based on phenotypic and ecological data [2]. However, the ability of adjusting cyanobacteria to fluctuating environmental conditions with morphological and biochemical modifications increases their diversity even at the intra-species level. On the other hand, this morphological

 $(\mathbf{\hat{P}})$ (cc) BY NC

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/by-nc/3.0/) which per-

mits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2021 Korean Society of Environmental Engineers

and biochemical complexity makes the identification and evaluation of their diversity more and more complicated. Therefore, morphological identification is not sufficient to make conclusions on the classification and the taxonomical state of a particular cyanobacteria strain. Higher variability of morphological features with fluctuating environmental conditions is the major problem of using it alone as the major criterion of cyanobacteria classification [3]. Therefore, molecular characterization is essential for the confirmation of the identity and their state in taxonomy.

Precise identification of cyanobacteria is thus essential in an accurate evaluation of cyanobacteria diversity. Identification of cyanobacteria which is solely based on the morphology would not be reliable as the morphological features are not static as they are modified with environmental fluctuations. This highlights the importance of using a polyphasic approach in the diversity evaluation. A combination of morphological and molecular charac-

[†] Corresponding author

Received January 26, 2020 Accepted April 28, 2020

Email: renuka.ra@nifs.ac.lk / renukaratnayake12@gmail.com Tel: +94 812232002 Fax: +94 812232131

terization can therefore be used as a polyphasic approach to evaluate the cyanobacteria diversity.

Sri Lanka is one of the major biodiversity rich representatives in the tropical region with different climatic zones in which many different ecosystems and vegetations are found. There is a huge variation of floral, faunal and microbial diversity among these zones due to remarkable variations in environmental conditions. This tropical continental island bears number of inland freshwater bodies including 103 rivers, over 250 water supply systems [4] and about 10,000 man-made lakes that periodically experience extreme dry conditions which lead to eutrophic conditions [5]. Favorable environmental conditions such as the atmospheric, water and soil temperature, humidity, solar irradiation, pH and precipitation allow not only animals and plants but also mostly undiscovered microbial communities to flourish. Diverse collection of ecosystems and water sources therefore implies the remarkable diversity of cyanobacteria communities among all other microbial communities in this region.

A broad range of fluctuating environmental conditions throughout the year and the presence of number of different ecosystems and water sources within a small region make Sri Lanka one of the best biodiversity hot spots in the tropical Asian region to study the diversity of cyanobacteria. However, the studies carried out so far, exploring the diversity, distribution and the identification of cyanobacteria in Sri Lanka are scarce. Recently few reports have been published on molecular characterization of cyanobacteria in Sri Lanka. Among them, the studies applying a polyphasic approach are even more limited. The cyanobacteria diversity in different water reservoirs, paddy fields, brackish water and tsunami affected areas in Sri Lanka has been studied using a polyphasic approach including the combination of light microscopy, cultures and the analysis of 16S rRNA gene sequences by Wanigatunge et al. [6]. Sethunge and Manage [4] have identified toxin producing, filter clogging, taste and odor forming nuisance algae and some physio-chemical parameters in some selected water bodies in Sri Lanka. Bloom formation of cyanobacteria along with dominance of Microcystis sp. was reported in 17 reservoirs of fresh water bodies in Sri Lanka by Javatissa et al. [7]. Genetic divergence, phylogenetic relationships and potential toxin producing cyanobacteria in reservoirs and well waters in Dry zone of Sri Lanka have been studied using 16S rRNA gene sequences by Livanage et al. [8]. This study confirms a rich diversity and divergence among toxin producing cyanobacteria in the water bodies in Dry zone in Sri Lanka [8]. Almost all the studies carried out in Sri Lanka so far have been focused on the occurrence of cyanobacteria and their toxic effects. The information on cyanobacterial culturing, biomass harvesting and the use of biomass for various industrial purposes have remained rare or scarce.

Cyanobacteria can provide promising solutions to most of the prevailing global issues. Nutritious food crisis is one such problem. Rapid growth of human population accelerates the depletion of arable lands for cultivation and leads a huge food crisis. Less growth requirements and high nutrient profiles of cyanobacteria make them promising food supplements and have already been used as food sources especially in Asia. *Spirulina* sp. [9] and *Aphanotheca sacrum* [10] had been added for their diet by local populations daily basis in some of the Asian countries.

Rapid depletion of protective Ozone layer due to higher amounts of harmful greenhouse gasses has caused harmful Ultra Violet (UV) rays to reach the Earth. Cancerous effects on eyes and skin caused by UV light have thus led to major globally prevailing health issues. With the higher demand for UV protective agents, synthetic UV protective agents were popularized. However, potential toxicity shown by these products has increased the demand for naturally available UV protective agents. Some herbal extracts play a major role in UV protection. Naturally occurring cyanobacteria is one of the best alternatives with high UV protective properties.

Toxin production is one of the well-known natural defense mechanisms in cyanobacteria. Under adverse environmental conditions, some of the strains are producing different types of cyano toxins. Value added potential of *C. komarekiana* highlights wide industrial applicability of the species thus the safety level of application and consumption should be further studied and confirmed. Therefore, their capability of toxin production and the levels of toxin production should be essentially evaluated before providing recommendations.

Therefore, the main objective of this study was to investigate the diversity of culturable cyanobacteria from freshwater bodies representing three major climatic zones; Dry zone, Intermediate zone and Wet zone in Sri Lanka, using a phylogenetic perspective. The specific objectives were (1) to isolate cyanobacterial strains followed by their morphological characterization (2) to investigate their phylogenetic relationships using the 16S rRNA gene sequence (3) to compare these sequences with other published sequences for deeper taxonomic resolution and (4) to identify and record new species with their value added potential and toxin producing capability.

These goals will improve our knowledge regarding the diversity, ecological distribution and industrial applicability of culturable cyanobacteria strains from Sri Lanka. We hypothesized that freshwater bodies in the tropical region Sri Lanka, harbor an unknown diversity of unicellular and filamentous cyanobacteria that have never been reported for this region.

2. Material and Methods

2.1. Study Area

Sampling was conducted from July 2014 to May 2015. Water samples were collected from freshwater bodies representing Dry zone, Intermediate zone and Wet zone of Sri Lanka including Jaffna peninsula as a major region in Dry zone of Sri Lanka. Sampling was carried out in triplicates, once in each location. Most of the reservoirs selected for sampling were man made and from dry zone of Sri Lanka.

2.2. Sample Collection and Preparation

Water was collected from the photic surface layer of the reservoir using a Ruttner sampler and kept in a plastic can till the sample was prepared for culturing (less than two hours). The photic zone of the water was determined by the Secchi disk disappearance technique. The coordinates of all the sampling locations were taken using Geographical Positioning System (GPS) receiver



Fig. 1. Sampling locations of freshwater bodies in Sri Lanka.

(Garmin, eTrex 30). Sampling locations are summarized in Fig. 1. From each sample 2 L volume was filtered through 20 μ m mesh size planktonic net. The retentiate was transferred into a screw cap plastic tube making the final volume into 25 mL by adding distilled water. Another sample was prepared by adding original water to the retentiate instead of distilled water. During sampling, physical parameters (water temperature, pH, and Secchi depth (cm)) of the freshwater bodies in three climatic zones were also measured.

2.3. Cyanobacteria Isolation and Culturing

Ten mL of the planktonic samples were transferred into 50 mL of BG11 [11], GO [12] and ASNIII [13] media for culturing. Ten mL of samples with 40 mL of medium were cultured in 100 mL conical flasks for laboratory cultures and 10 L aspirator bottles with one third filled with media were used for semi mass culturing which were kept at shaking (0 rpm, 200 rpm and 300 rpm) in biological growth chamber under fluorescent light (1000 lux, 1500 lux and 2000 lux light intensity) with constant illumination. Four different pH (7, 7.5, 8, and 8.5) values were selected and growth pH range was optimized based on the average pH range shown by six representative water samples collected from three major climatic zones. The pH of the medium was adjusted using 10% Sodium Hydroxide and 10% Hydrochloric Acid. Number of days taken by cultures in each medium to show their bluish green color growth was recorded. Once growth was observed, 100 μ L of samples

were sub cultured into petri-dishes by the spread plate method under a laminar air flow cabinet [14] on agar plates containing the same media solidified with 1.5% (w/v) bacteriological agar.

Frequent sub culturing was carried out to isolate single colonies from mixed cultures of cyanobacteria. Repeated streaking and spreading on to the solid medium and microscopic observations were performed until pure monocultures were obtained. Vortexing was used to separate the filamentous cyanobacteria when they were in mixtures. For this, one mL of the sample along with some pieces of glass beads was poured into a micro-centrifuge tube and vortexed. In case of plate culture, algal colonies were picked up using a sharp needle and diluted with water in micro-centrifuge tubes prior to vortexing. Subcultures were obtained by inoculating 100 μ L of the vortexed sample on to a petri plate. The purity of the culture was confirmed by repeated sub culturing in plate and regular observation under the microscope [15]. After isolation, all the cyanobacterial strains were cultured and maintained in 250 mL Erlenmeyer flasks filled with 200 mL of BG-11 medium. They were also cryopreserved in a -80°C freezer with cryoprotective agents.

2.4. Morphological Identification of Isolated Cyanobacteria Strains

Morphological identification of the isolated strains was conducted using a compound microscope (Euromex BioBlue.Lab BB. 1153-PLi) equipped with the image system, Image Focus 4 ver.2.9 and the classification system was devised by Desikachery [16] as well as recently published literature [17].

2.5. Molecular Analysis of Selected Cyanobacteria Isolates

Based on the efficiency in biomass production, a total number of 31 isolates were selected for molecular identification with the purpose of evaluating their value added potential for different industrial applications.

2.5.1. DNA extraction and 16S rRNA gene region amplification and sequencing

Total genomic DNA was extracted from the cultured cyanobacterial cells using the method of Smoker and Barnum [18] with slight modifications. The 16S rRNA gene region was PCR amplified in a Techine TC 3000 DNA Thermal Cycler using the primer sets; CYA106F (approx. 675 bp in length) and CYA359F (approx. 450 bp in length) as forward primers and CYA781Ra and CYA781Rb as reverse primers [19].

The cycling profile for the amplification of 16S rRNA gene region included an initial denaturation of template DNA at 94°C for 5 min, followed by 40 cycles of 94°C for 1 min, annealing at 60°C for 1 min, elongation at 72°C for 1 min and a final elongation at 72°C for 15 min. The amplified PCR products were purified using a Wizard SV gel and PCR clean up system PCR purification kit (Promega Corporation, USA) as recommended by the manufacturer. Sequencing of 16S rRNA gene was carried out at Macrogen, South Korea using ABI 3730XL sequencers with amplified samples of respective forward and reverse primers.

2.5.2. BLAST analysis and phylogenetic tree of 16S rRNA gene The organisms were identified at a molecular level using online Blast software available in NCBI website. The partial sequences obtained in this study were compared with existing sequences for cyanobacteria in NCBI database. The nucleotide sequences generated in this study and the reference sequences retrieved from GenBank were multiple-aligned using CLUSTAL W, Version 1.7. The alignment was corrected manually and converted to a distance matrix. The distance matrix was converted to a phylogenetic tree using the Neighbour-joining (NJ) algorithm [20] of MEGA 5.05; [21] with multiple substitutions corrected and positions with gaps excluded. The 16S rRNA gene sequences determined for the freshwater cyanobacterial strains were deposited in the GenBank database via BankIt submission tool under the accession numbers: KX962074 - KX962104.

2.5.3. Identification and confirmation of the identity of new species *C. komarekiana*

Depending on the DNA sequencing results, two strains; U08 and U41 were identified as *C. komarekiana* whose presence in Sri Lanka was identified and reported for the first time. The identity was further confirmed with reproducible DNA sequencing results and morphological characteristics. Furthermore, the value added potential of *C. komarekiana* was determined as follows.

2.6. Investigation of Value Added Potential of C. komarekiana

2.6.1. Analysis of nutrient content in C. komarekiana

Mass culturing of *C. komarekiana* was carried out in 100 L size fish tanks under greenhouse environmental conditions with natural light and temperature. Media was prepared at half strength of respective BG-11 at pH 7.5 along with aeration (Risheng RS 2800). Centrifugation, filtering and flocculation techniques (by adjusting pH) were used to harvest the biomass and dry biomass was obtained and made it into a fine powder for further analysis.

2.6.1.1. Total carbohydrate content of C. komarekiana

The Dubois method [22] was used to measure the total carbohydrate content of the fresh dry biomass. The absorbance of the samples was measured at a wavelength of 490 nm using a microplate reader in a spectrophotometer (UV-2450, Shimadzu).

2.6.1.2. Total protein content of C. komarekiana

Lowry method [23] where the absorbance of the sample was measured at a wavelength of 750 nm in spectrophotometer, was used to measure the total protein content of the fresh biomass. The spectrophotometric absorbance was converted to protein concentration using a calibration curve established with Bovin Serum Albumin.

2.6.1.3. Macro and micro mineral contents of *C. komarekiana* 0.1 g of dried cyanobacteria biomass was measured and placed in the microwave digester and 3 mL of 69% HNO₃ was applied and digested them for 25 min. After cooling, the digested volume was transferred to 10 mL volumetric flasks and volumed up to 10 mL with ultra-pure water. Then the diluted volume was filtered using 0.45 μ m filter papers. The resulted filtrate was again filtered using 0.45 μ m membrane filters and the final filtrate was used for the analysis by using Inductively Coupled Plasma Optical Emission Spectrometry (ICP-OES). Among macro minerals, Ca was analyzed by Aggressive Digestion Method using Atomic Absorption Spectrophotometer (AAS)

2.6.2. Measurement of Sun Protection Factor (SPF)

Sun Protection Factor (SPF) was determined by the method described by Dutra et al. [24] followed by the application of Mansur equation shown in Eq. (1) [25]. The absorption data were obtained in the range of 290 to 320, every 5 nm, and 3 determinations were made at each point.

$$SPF_{in \ vitro} = CF \times \sum_{290}^{320} EE(\lambda) \times I(\lambda) \times abs(\lambda)$$
(1)

Where: EE - Erythemal effect spectrum; I - Solar intensity spectrum; Abs- Absorbance of sunscreen product; CF - correction factor (= 10). SPF value was determined by UV spectrophotometer [25].

2.7. Determination of Toxin Producing Capability of *C. komarekiana*

Toxin analysis and quantification were carried out by using ELISA (Enzyme-Linked Immunosorbent Assay) at the Department of Zoology, University of Sri Jayawardenapura, Sri Lanka. Mainly, the toxins; Microcystin-LR and Cylindrospermopsin were analyzed and quantified.

3. Results and Discussion

3.1. Cyanobacteria isolation and morphological identification

In the present study a total of 74 uni-algal cultures were obtained and these were distributed as 52, 16 and 6 from reservoirs of the Dry, Wet and Intermediate zones respectively (Table 1). The morphological characteristics of cyanobacteria which were observed during the present study are summarized in Table S1. Traditionally, the classification of cyanobacteria has been based on morphological characters such as trichome width, cell size, division planes, shape, arrangement, pigmentation and the presence of characters such as gas vacuoles and a sheath [26, 27]. However, some diagnostic features such as gas vacuoles or akinetes can show variations with different environmental or growth conditions and even be lost during cultivation [28, 29]. Such limitations of phenotypic characters have highlighted the requirement for more reliable methods and promoted molecular approaches in cyanobacterial taxonomy including DNA base composition [30, 31], DNA hybridizations [32], gene sequencing [19] and PCR fingerprinting [33, 34].

Compared to other climatic zones, Wet zone receives relatively high mean annual rainfall over 2,500 mm without pronounced dry periods. The Dry zone receives a mean annual rainfall of less than 1,750 mm with a distinct dry season from May to September. The Intermediate zone receives a mean annual rainfall between 1,750 to 2,500 mm with a short and less prominent dry season [35]. According to the present study, the highest water temperature (31.35 \pm 1.5°C), pH (8.20 \pm 0.68) and Secchi depth (75.09 \pm 26.31 cm) were recorded in Dry zone whereas all these parameters were recorded the lowest in the Wet zone (Table S2). This could be

Agro ecological Zone	the morphological characteristics		
	Oscillatoria sp.		
	Phormidium sp.		
	Pseudoanabaena sp.		
	<i>Gleocapsa</i> sp.		
	Planktolyngbya sp.		
	Microcystis sp.		
	Limnothrix sp.		
	Microcoleus sp.		
Dry zone	Aphanothece sp.		
	Synechococcus sp.		
	Lyngbya sp.		
	Planktothrix sp.		
	Chroococcidiopsis sp.		
	Anabaena sp.		
	Nostoc sp.		
	Microchaete sp.		
	Anabaena sp.		
	Plectonema sp.		
	<i>Lyngbya</i> sp.		
	Calothrix sp.		
	Limnothrix sp.		
	Chroococcus sp.		
Wet zone	Phormidium sp.		
	Chroococcidiopsis sp.		
	Dermocarpa sp.		
	Leptolyngbya sp.		
	Nostoc sp.		
	Oscillatoria sp.		
	Synechococcus sp.		
	Microcystis sp.		
	Oscillatoria sp.		
	Phormidium sp.		
Intermediate	Microcystis sp.		
zone	Synechococcus sp.		
	Chroococcales sp.		
	<i>Plectonema</i> sp.		

 Table 1. Identification of Cyanobacteria Uni Algal Cultures Based on the Morphological Characterization

due to higher rainfall with shorter dry season in the Wet zone which directly influences the water temperature. On the contrary, the lowest rainfall in the Dry zone of the country prolongs the dry season which also influences the water temperature. Favorable environmental conditions such as sufficient amounts of sunlight, water temperature, pH and nutrient rich environment could cause higher diversity of cyanobacteria in Dry zone.

Though 45 genera are reported so far from this tropical region, Sri Lanka [36], a recent study carried out by Wanigatunge et al. [6] reported 26 genera out of them. Another study reports about 40 species of cyanobacteria belonging to 24 genera from Sri Lankan reservoirs [37]. Of these, except for *Microcystis aeruginosa*, other species are either rare or occur only in small numbers [37]. More recent reports on the occurrence of freshwater cyanobacteria give evidence of *Cylindrospermopsis* as a dominant form in addition to *Microcystis* in several reservoirs surveyed [5, 8, 38]. However in the present study *Cylindrospermopsis* was not observed among the genera isolated and cultured.

3.2. Molecular Identification-The 16S rRNA Region Analysis

3.2.1. Sequencing of 16S rRNA region analysis

The partial sequences obtained in this study were compared with existing sequences for cyanobacteria in NCBI database. All the sequences in the present study showed \geq 95% similarity with either query coverage or identity or both with existing data. Hence, they were identified with the closest matching data (Table 2). The selection of 31 isolates for molecular identification was based on efficient biomass production.

In the present study, though 74 isolates have been reported based on morphology, molecular identification was carried out only for selected strains which were efficient in biomass production to check their value added potential to be utilized in different industries.

3.2.2. First record on the occurrence of *Cephalothrix komarekiana* in Sri Lanka

Molecular identification results report the availability of some rare isolates such as Alkalinema pantanalense (U4), Geitlerinema sp. (U36), Westiellopsis prolifica (U58) which were reported in 1986 from this tropical island [36] or rarely been observed thereafter in Sri Lanka. Among them, the novel species Cephalothrix komarekiana was identified for the first time from freshwaters of Sri Lanka (Fig. 2). With morphological and molecular characterization, two isolates; U8 and U41 were identified as Cephalothrix komarekiana. All these two water samples were collected from the freshwater bodies named as Ariva Kulam (U8) and Pollu Kulam (U41) which are located in the northern part of the country, Jaffna peninsula in Sri Lanka. Jaffna peninsula is in the northern most part of Sri Lanka and based on annual average rainfall it covers a major area in Dry zone of Sri Lanka (5° 54' N -9° 52' N and 79° 39' E - 81° 53' E). The average mean temperature of the study area varies in between 25.0 to 27.5°C while annual cumulative rainfall is reported in between 1000 to 1500 mm [39]. Rainfall of the study area is seasonal and mainly governed by the North East Monsoon. According to the water quality reports, the freshwater of Jaffna is alkaline in character with high salinity.

The 16S rRNA gene sequence showed 99% of query cover and 97% of identity against the best hit available in NCBI public database (Table 2) for the isolate U8. The query cover and identity against the best hit were 100% for the isolate U41 (Table 2).

ID	Molecular identification	Query cover	Identity	Genbank accession no
U-1	Leptolyngbya sp.AW1	99%	99%	KX962074
U-2	Phormidium sp.AW2	100%	99%	KX962075
U-3	Limnothrix sp.KW3	100%	100%	KX962076
U-4	Alkalinema pantanalense CW4	100%	99%	KX962077
U-6	Anabaena sp.CW6	95%	98%	KX962078
U-7	Phormidium sp.KW7	100%	99%	KX962079
U-8	Cephalothrix komarekiana AK8	99%	97%	KX962104
U-9	*Limnothrix sp.HW9	99%	89%	KX962080
U-10	Synechococcus sp.KT10	99%	98%	KX962081
U-12	Synechocystis sp.GL12	100%	100%	KX962082
U-13	Chroococcidiopsis sp.UW13	100%	91%	KX962083
U-15	Calothrix sp.KK15	97%	94%	KX962084
U-16	Chroococcidiopsis sp.GL16	99%	100%	KX962085
U-22	*Cephalothrix sp.MK22	100%	91%	KX962086
U-27	Phormidium sp.KL27	99%	98%	KX962087
U-30	*Limnothrix sp.UW30	92%	88%	KX962088
U-31	Calothrix sp.KE31	92%	98%	KX962089
U-33	Limnothrix sp.UW33	100%	96%	KX962090
U-36	Geitlerinema sp.KK36	100%	100%	KX962091
U-38	Synechococcus sp.WW38	100%	99%	KX962092
U-40	Oscillatoriales sp.PK40	99%	92%	KX962093
U-41	Cephalothrix komarekiana PK41	100%	100%	KX962094
U-42	Synechocystis sp.MW42	99%	98%	KX962095
U-49	Nostoc sp.AW49	99%	99%	KX962096
U-50	Calothrix sp.AW50	100%	99%	KX962097
U-55	Oscillatoriales sp.DL55	99%	92%	KX962098
U-57	Plectonema sp.KT57	100%	96%	KX962099
U-58	Westiellopsis prolificaKL58	100%	99%	KX962100
U-63	Limnothrix sp.MK63	100%	98%	KX962101
U-67	Limnothrix sp.NW67	100%	100%	KX962102
U-106	Anabaena sp.CW106	95%	99%	KX962103

Table 2. Molecular Identification of Cyanobacteria in Reference with BLAST

(*Indicates the strains which have given a query cover or identity less than 92%. The genera were assigned based on the Blast analysis).



Fig. 2. Microscopic images of Cephalothrix komarekiana U41 and U8.

3.2.3. Confirmation of the identity of the new species, *Cephalothrix* komarekiana

Polyphasic approach where the combination of morphological and molecular identification approaches were used, helped in further confirmation of the identity of the new species. DNA sequencing results were highly reproducible to confirm the identity of the isolate U41 as *Cephalothrix komarekiana*. Both percentage identity and query cover were considered for identification of this strain up to species level. Reproducible results showed 100% query cover and 99.43% percentage identity for U41 (Table 3) confirming the identity of the isolate as *Cephalothrix komarekiana*.

In previous studies, this novel species was first reported from tropical South American region [40, 41] and a temperate region in Korea [42]. The first report stated that the species was existed in extremely alkaline water and freshwater aquarium in Brazil in 2015 [40] and then from freshwater of Amazonian rivers in the same South American region, Brazil in 2017 [41]. The same species was recorded from brackish waters in Korea in 2018 [42]. However, the identified strains in Sri Lanka were present in freshwater reservoirs used for irrigation purposes. This study has reported the first record of *Cephalothrix komarekiana* from freshwater reservoirs of Sri Lanka; a country in the Tropical Asia with different environmental conditions.

3.3. Construction of the Phylogenetic Tree with the DNA Sequences Obtained in the Study

According to the constructed phylogenetic tree, it was observed that the isolate U9, U22, U30 and U31 had lower than 95% query coverage or identity. However, the phylogenetic tree with the sequences obtained in the present study for those genera along with previously sequenced sequences obtained from GenBank showed all isolates under the same branch but in different clusters (Fig. 3). Though the isolate U22 was morphologically similar to the two isolates *Cephalothrix komarekiana* PK41 and *Cephalothrix komarekiana* AK8, partial sequencing indicated only 91% sequence similarity to the two isolates. By subjecting these three isolates for the total sequencing of the 16S rRNA gene, the confirmation of the species could be achieved.

3.4. Value Added Potential of C. komarekiana

The present study also reports for the first time on the value added potential of *Cephalothrix komarekiana* present in freshwater bodies in Sri Lanka. The evaluation of value added potential of this newly recorded species will provide a new insight to the applied research and related industrial applications.

3.4.1. Analysis of nutrient contents in C. komarekiana

Total carbohydrate content of the strain of C. komarekiana is 33.23% (Table 4). This value is considerably higher with comparison to the records on proximate available carbohydrate contents present in three microalgae strains; Spirulina plantensis, Chlorella vulgaris and Isochrisis galbana [43]. However, Spirulina plantensis was recorded with a higher average crude protein content of 63.0% [43] whereas the above strain of C. komarekiana showed an average total protein content of 28.93% (Table 4). The fatty acid content of C. komarekiana was recorded in between 12-14% (Table 4) and this strain was among the cyanobacteria strains with considerably higher fatty acid contents. Considerably high major nutrient contents of C. komarekiana suggest their suitability as an alternative food supplement. Furthermore, their nutrient profile for important macro and micro mineral contents will provide a broad picture on their effectiveness and suitability in developing this species as a food supplement.

Nutrient profiles of macro and micro nutrients show that C. komarekiana is rich with higher amounts of macro nutrients such as Mg, K and Ca. Among the tested macro nutrients, Ca was the highest that was analyzed by AAS. Mg was the second highest followed by K (Table 5). Ca was reported as a rich mineral in Spiruling sp. which is already popular as a food source in the world [44]. The reported concentration of Ca in Spirulina sp. was in the range of 1,300-14,000 ppm and it was comparable to the quantity found in milk [44]. Similarly, in this study, the Ca content in C. komarekiana was reported as 4258 ppm. Therefore, C. komarekiana can be identified as a rich source of Ca. As well as Spirulina sp., C. komarekiana therefore becomes a suitable nutrient supplement for vegetarians. Compared to other macro elements, Ca is required by the cellular systems in large amounts as they are essential for cellular communication and modulation including cell movement, muscle contraction, nerve transmission, glandular secretion, blood clotting and cell division. As a Ca rich cyanobacterium, C. komarekiana shows its value added potential to be developed as a low cost, alternative supplement for Ca. Not only Ca but also higher amounts of Mg and K in C. komarekiana suggest its potential to be utilized in food industry and in different medical applications. Iron was in the greatest quantity followed by Mn, Zn, Cu, Co and Ni among all micro nutrients (Table 5). C. komarekiana showed a higher Cu content (4.5 ppm) compared to Spirulina sp. (1.9 ppm) [45]. Mn content present in C. komarekiana (30.3 ppm) is similar to the Mn content in Spirulina sp. (31.2 ppm) [45]. However, Zn (5.7 ppm) and Mg (693.0 ppm) contents in C. komarekiana were lower compared to Spirulina sp. (32.0 ppm and 2120.0 ppm respectively) [45]. Heavy metal quantities of Cd, Cr, Pb, As and Sr in C. komarekiana were in the safe consumption level recommended

Table 3. Confirmation of Molecular Identification of the New Species in Reference with BLAST

ID	Molecular Identification	Query cover	Identity	Genbank Accession No
U41	Cephalothrix komarekiana SAG 75.79	100%	99.43%	MN908259

Table 4. Major Nutrient Contents (as a percentage of dry biomass) in the New Strain of C. komarekiana

Total Carbohydrate content (%)	Total Protein content (%)	Fatty acid content (% dry biomass)
33.23%	28.93%	12-14%



Fig. 3. Phylogenetic tree for the isolates with lower than 95% query coverage and identity in comparison with the sequences obtained in the present study (starting with the symbol 'U') along with the previously sequenced sequences obtained from GenBank database (starting with the symbol 'seq').

 Table 5. Macro and Micro Mineral Contents (in ppb or ppm) in C. komarekiana

Macro/ Micro minerals	Concentration [ppb or ppm)
Cd [ppb]	7.63
Ni [ppb]	31.97
Zn [ppb]	5744.47
Mn [ppb]	30314.80
Cr [ppb]	158.64
Pb [ppb]	40.98
Co [ppb]	888.87
Cu [ppb]	4514.21
As [ppb]	11.67
Fe [ppm]	146.85
Mg [ppm]	693.07
Ca [ppm]	4258.00
K [ppm]	374.90
Sr [ppb]	197.70

by the World Health Organization. These concentrations in *C. ko-marekiana* were also reported in negligible amounts or in very low concentrations and always lower than the reported levels in *Spirulina* sp. [45]. It further confirms their level of safety in utilizing them as mineral supplements in food and pharmaceutical industries. As a mineral rich cyanobacterium, *C. komarekiana* would be another promising natural alternative which can be developed as a food supplement for the improvement of both global public health and economy.

3.4.2. Measurement of Sun Protection Factor (SPF) in *C. komarekiana C. komarekiana*: SPF value: 2.37 ± 0.755

According to the above results, the reported SPF value for *C. komarekiana* is 2.37 ± 0.755 . Compared to the SPF values obtained for some of the herbal extracts namely; Aloe vera, Carrot, Cucumber and Watermelon by Malsawmtluangi et al. [46], *C. komarekiana* has shown considerably higher SPF value. Therefore, it suggests the effectiveness of *C. komarekiana* as a promising botanical agent to be utilized and developed as a UV protective product in cosmetic industry. Higher photosynthetic ability and growth rate, less nutrient

Suitable nutrient profile and efficient and high UV protective properties suggest the value added potential of *C. komarekiana* and provide a clear insight on possible industrial applications. However, some cyanobacteria strains are toxin producing. Therefore, recommendation of the species for possible industrial applications is still one step away and requires further confirmation of their level of risk for applications.

3.5. Toxin Analysis of C. komarekiana

Toxin producing capability of C. komarekiang for Microcystin-LR and Cylindrospermopsin was investigated. Total Microcystin-LR for C. komarekiana was $0.63 \pm 0.045 \ \mu g/L$. The detected concentration of total Cylindrospermops in for the strain was 0.06 ± 0.085 μ g/L. According to the World Health Organization (WHO)'s and the Oregon Health Authority (OHA)'s guidance on cyanotoxins, a provisional guideline value of 1.0 μ g/L has been adopted for Microcystin-LR. The above guideline value is applicable for all cell bound and extracellular microcystins [47, 48]. Provisional lifetime guideline value set by WHO for Cylindrospomopsin is 0.7 μ g/L and short-term guideline value is 3.0 μ g/L [47]. The OHA's provisional guideline value for Cylindrospomopsin is $1.0 \ \mu g/L$ [48]. All these values have been calculated considering average adult body weight of 60 kg as a standard [47, 48]. With comparison to all these standard values, all the produced toxin concentrations are below the risk level for human health. Therefore, these values further confirm the suitability of using C. komarekiana as a food supplement, effective natural sun protective agent and in other industrial applications.

4. Conclusions

The present study was carried out to investigate the diversity of culturable cyanobacteria from freshwater bodies in the tropical Asian region Sri Lanka, with a phylogenetic perspective. A total of 74 uni-algal cultures were obtained and these were distributed as 52, 16 and 6 from reservoirs of the dry, wet and intermediate zones respectively. Among them, C. komarekiana was recorded for the first time from tropical region; Sri Lanka with the availability of some other rare isolates such as Alkalinema pantanalense, Geitlerinema sp., and Westiellopsis prolifica. Rich nutrient profiles of total carbohydrates, total proteins, fatty acids and macro and micro elements showed its capability to be developed as natural, safe alternative food and mineral supplements. Rich Ca profile in C. komarekiana suggests its potential utilization separately as a Ca supplement which would be useful in treating complications caused by Ca deficiencies. C. komarekiana has also been proved to be an efficient natural UV protective agent. Suitable nutrient profiles and efficient UV protective properties provide a clear insight on a wide range of possible industrial applications of C. komarekiana. Their ability and concentrations of toxin production further confirmed the safety level of utilization of this species for the improvement of public health. Higher photosynthetic ability and growth rate, less nutrient and land area requirement, capability of mass production with less capital investment and most importantly zero environment pollution nominate this species as one of the best natural agents which can be utilized in finding eco-friendly solutions for many of the global issues.

Nomenclature

- $\Sigma \quad \text{Summation of}....$
- Λ Wavelength

Acknowledgment

The authors would like to thank Prof. M.M. Pathmalal from Department of Zoology, University of Sri Jayawardenapura for providing facilities for Toxin analysis.

Author Contributions

M.F. (Assistant Professor) conducted all the laboratory work and experiments and data analysis starting from sampling to identification of new species. R.W.T.M.R.T.K. (M.Phil student) conducted experiments and data analysis to confirm the identity of the new species and to analyze the value added potential of the new species and manuscript preparation. K.L.W. (Professor) supervised the above research work, reviewed and corrected the manuscript. D.N. (Associate Research Professor) supervised the research work related to molecular analysis, reviewed and corrected the manuscript. R.R. (Senior Research Fellow) supervised the above research work, reviewed and corrected the manuscript.

References

- Oren A. Cyanobacterial systematics and nomenclature as featured in the International Bulletin of Bacteriological Nomenclature and Taxonomy/International Journal of Systematic Bacteriology/ International Journal of Systematic and Evolutionary Microbiology. Int. J. Syst. Evol. Microbiol. 2011;61(1):10-15.
- Komárek J. Recent changes (2008) in cyanobacteria taxonomy based on a combination of molecular background with phenotype and ecological consequences (genus and species concept). *Hydrobiologia*. 2010;639(1):245-259.
- Palinska KA, Liesack W, Rhiel E, Krumbein WE. Phenotype variability of identical genotypes: the need for a combined approach in cyanobacterial taxonomy demonstrated on *Merismopedia*-like isolates. *Arch. Microbiol.* 1996;166(4):224-233.
- Sethunge S, Manage PM. Nuisance algae in water supply projects in Sri Lanka. International Conference on Sustainable Built Environment; 13-14 December 2010; Kandy, Sri Lanka. p. 62-70.
- Perera MBU, Yatigammana SK, Kulasooriya SA. Prevalence of toxigenic cyanobacteria in different climatic zones of Sri Lanka. International Symposium on Water Quality and Human Health: Challenges Ahead; 2012; Postgraduate Institute of

Science, University of Peradeniya, Sri Lanka.

- 6. Wanigatunge RP, Magana-Arachchi DN, Chandrasekharan NV, Kulasooriya SA. Genetic diversity and molecular phylogeny of cyanobacteria from Sri Lanka based on 16S rRNA gene. *Environ. Eng. Res.* 2014;19(4):317-329.
- Jayatissa L, Silva E, McElhiney J, Lawton L. Occurrence of toxigenic cyanobacterial blooms in freshwaters of Sri Lanka. Syst. Appl. Microbiol. 2006;29(2):156-164.
- Liyanage HM, Arachchi DNM, Chandrasekaran NV. Genetic divergence among toxic and non-toxic cyanobacteria of the dry zone of Sri Lanka. *SpringerPlus*. 2016;5(1):2026.
- 9. Abdulqader G, Barsanti L, Tredici MR. Harvest of *Arthrospira platensis* from Lake Kossorom (Chad) and its household usage among the Kanembu. *J. Appl. Phycol.* 2000;12:493-498.
- 10. Fujishiro T, Ogawa T, Matsuoka M, Nagahama K, Takeshima Y, Hagiwara H. Establishment of a pure culture of the hitherto uncultured unicellular cyanobacterium *Aphanothece sacrum*, and phylogenetic position of the organism. *Appl. Environ. Microbiol.* 2004;70(6):3338-3345.
- Stanier R, Kunisawa R, Mandel M, Cohen-Bazire G. Purification and properties of unicellular blue-green algae (order chroococcales). *Bacteriol. Rev.* 1971;35(2):171.
- Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. J. Gen. Microbiol. 1979;111(1):1-61.
- Rippka R, Waterbury JB, Stanier RY. Isolation and purification of cyanobacteria: Some general principles, The prokaryotes. *Springer.* 1981;212-220.
- Pulz O, Gross W. Valuable products from biotechnology of microalgae. Appl. Microbiol. Biotechnol. 2004;65(6):635-648.
- Abou-Shanab AR, Hwang JH, Cho Y, Min B, Jeon BH. Characterization of microalgal species isolated from fresh water bodies as a potential source for biodiesel production. *Appl. Energy.* 2011;88(10):3300-3306.
- Desikachary T. Cyanophyta. 8th ed. New Delhi: Indian Council of Agricultural Research; 1959. p. 686.
- Souza DT, Genuário DB, Silva FSP, et al. Analysis of bacterial composition in marine sponges reveals the influence of host phylogeny and environment. *FEMS Microbiol. Ecol.* 2017;93(1).
- Smoker JA, Barnum SR. Rapid small-scale DNA isolation from filamentous cyanobacteria. *FEMS Microbiol. Lett.* 1988;56(1): 119-122.
- Nübel U, Garcia-Pichel F, Muyzer G. PCR primers to amplify 16S rRNA genes from cyanobacteria. *Appl. Environ. Microbiol.* 1997;63(8):3327-3332.
- Saitou N, Nei M. The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 1987;4(4):406-425.
- Tamura K, Dudley J, Nei M, Kumar S. Mega4: Molecular evolutionary genetics analysis (mega) software version 4.0. *Mol. Biol. Evol.* 2007;24(8):1596-1599.
- Dubois M, Gilles KA, Hamilton JK, Rebers PT, Smith F. Colorimetric method for determination of sugars and related substances. *Anal. Chem.* 1956;28(3):350-356.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. J. boil. Chem. 1951;193(1): 265-275.

- Dutra EA, Oliveira DAGDC, Kedor-Hackmann ERM, Santoro MIRM. Determination of sun protection factor (SPF) of sunscreens by ultraviolet spectrophotometry. *Braz. J. Pharm. Sci.* 2004;40(3):381-385.
- Mansur JDS, Breder MNR, Mansur MCDA, Azulay RD. Determination of the sun protection factor by spectrophotometry. *An. Bras. Dermatol.* 1986;61(3):121-124.
- Komarek J, Anagnostidis K. Modern approach to the classification system of cyanophytes. 2-chroococcales. Arch. Hydrobiol. Suppl. 1986;73(2):157-226.
- Komárek J, Anagnostidis K. Modern approach to the classification system of cyanophytes Nostocales. *Algol. Stud.* 1989; 247-345.
- Lyra C, Suomalainen S, Gugger M, et al. Molecular characterization of planktic cyanobacteria of anabaena, aphanizomenon, microcystis and planktothrix genera. *Int. J. Syst. Evol. Microbiol.* 2001;51(2):513-526.
- Rudi K, Skulberg OM, Larsen F, Jakobsen KS. Strain characterization and classification of oxyphotobacteria in clone cultures on the basis of 16S rRNA sequences from the variable regions v6, v7, and v8. Appl. Environ. Microbiol. 1997;63(7):2593-2599.
- Kaneko T, Nakamura Y, Wolk CP, et al. Complete genomic sequence of the filamentous nitrogen-fixing cyanobacterium anabaena sp. Strain pcc 7120. DNA Res. 2001;8(5):205-213.
- 31. Kaneko T, Sato S, Kotani H, et al. Sequence analysis of the genome of the unicellular cyanobacterium *Synechocystis* sp. Strain pcc6803. Ii. Sequence determination of the entire genome and assignment of potential protein-coding regions. *DNA Res.* 1996;3(3):109-136.
- Kondo R, Yoshida T, Yuki Y, Hiroishi S. DNA-DNA reassociation among a bloom-forming cyanobacterial genus, *microcystis. Int.* J. Syst. Evol. Micr. 2000;50(2):767-770.
- Rasmussen U, Svenning MM. Fingerprinting of cyanobacteria based on pcr with primers derived from short and long tandemly repeated repetitive sequences. *Appl. Environ. Microbiol.* 1998;64(1):265-272.
- Versalovic J, Koeuth T, Lupski R. Distribution of repetitive DNA sequences in eubacteria and application to fingerprinting of bacterial enomes. *Nucleic Acids Res.* 1991;19(24):6823-6831.
- Premaratne S, Premalal GGC. Country pasture/forage resource profiles: Sri Lanka. Rome: Food and Agriculture Organization; 2006. p. 1-21.
- Abeywickrama B. The genera of the fresh water algae of Sri Lanka, man and the biosphere national committee for Sri Lanka. Special Publication (6), part 1–UNESCO. 1986.
- Silva E, Wijeyaratne M. The occurrence of cyanobacteria in the reservoir of the mahaweli river basin in Sri Lanka. Sri Lanka J. Aquat. Sci. 1999;4:51-60.
- Hossain MF, Ratnayake R, Kumara KW. Isolation purification and culturing of cyanobacteria and microalgae towards biodiesel production: Current status. *Am. J. Biomass Bioenergy*. 2016;5(3):111-129.
- Sri Lanka. Minindoru Departamentuva. National atlas of Sri Lanka. 2nd ed. Colombo: Survey Department of Sri Lanka; 2007.
- Malone CFS, Rigonato CF, Laughinghouse IV, et al. Cephalothrix gen. Nov. (cyanobacteria): Towards an intraspecific phylogenetic evaluation by multilocus analyses. *Int. J. Syst. Evol.*

Microbiol. 2015;65(9):2993-3007.

- 41. Genuario DB, Vaz, MGMV de Melo IS. Phylogenetic insights into the diversity of homocytous cyanobacteria from Amazonian rivers. *Mol. Phylogenet. Evol.* 2017;116:120-135.
- 42. Yim BC, Kim JH, Yoon SR, Lee NJ, Lee OM. A Study of Seven Unrecorded Species of Benthic and Subaerophytic Cyanobacteria (Cyanophyceae, Cyanophyta) in Korea. *Korean J. Environ. Biol.* 2018;36(3):291-298.
- Tokuşoglu Ö. Biomass nutrient profiles of three microalgae: Spirulina platensis, Chlorella vulgaris, and Isochrisis galbana. J. Food Sci. 2003;68(4):1144-1148.
- 44. Falquet J. The nutritional aspects of Spirulina [Internet]. Antenna Foundation; c2020 [cited 25 April 2020]. Available from: https://www.antenna.Ch/wp-content/uploads/2017/03/ AspectNut_UK.pdf.
- 45. Alobwede E, Leake JR, Pandhal J. Circular economy fertilization: Testing micro and macro algal species as soil improvers and nutrient sources for crop production in greenhouse and field conditions. *Geoderma* 2019;334:113-123.
- Malsawmtluangi C, Nath DK, Jamatia I, Zarzoliana E, Pachuau L. Determination of Sun Protection Factor (SPF) number of some aqueous herbal extracts. J. Appl. Pharm. Sci. 2013;3(9):15.
- Chorus I, Bartram J., eds. Toxic Cyanobacteria in Water: A guide to their public health consequences, monitoring and management. London: CRC Press; 1999.
- 48. Farrer D, Counter M, Hillwig R, Cude C. Health-based cyanotoxin guideline values allow for cyanotoxin-based monitoring and efficient public health response to cyanobacterial blooms. *Toxins.* 2015;7(2):457-477.