



Article

Household Air Pollution from Biomass Fuel for Cooking and Adverse Fetal Growth Outcomes in Rural Sri Lanka

Alicia Vakalopoulos ¹, Shyamali C. Dharmage ^{1,*} , Samath Dharmaratne ^{2,3}, Pasan Jayasinghe ⁴ , Olivia Lall ¹, Isabella Ambrose ¹, Rohan Weerasooriya ⁵, Dinh S. Bui ¹ , Duminda Yasaratne ⁶, Jane Heyworth ⁷ and Gayan Bowatte ^{1,5,8}

- ¹ Allergy and Lung Health Unit, Melbourne School of Population and Global Health, University of Melbourne, Carlton, VIC 3053, Australia; avakalopoulo@student.unimelb.edu.au (A.V.); olall@student.unimelb.edu.au (O.L.); iiambrose@student.unimelb.edu.au (I.A.); dinh.bui@unimelb.edu.au (D.S.B.); gayan.bowatte@unimelb.edu.au (G.B.)
 - ² Department of Community Medicine, Faculty of Medicine, University of Peradeniya, Peradeniya 20400, Sri Lanka; samath.dharmaratne@mbbs.md
 - ³ Department of Health Metrics Sciences, Institute for Health Metrics and Evaluation, School of Medicine, University of Washington, Seattle, WA 98195, USA
 - ⁴ Department of Health Services, Central Province, Kandy 20000, Sri Lanka; pasan978@gmail.com
 - ⁵ National Institute of Fundamental Studies, Hantana Road, Kandy 20000, Sri Lanka; rohan.we@nifs.ac.lk
 - ⁶ Department of Medicine, Faculty of Medicine, University of Peradeniya, Peradeniya 20400, Sri Lanka; yasaratne@yahoo.com
 - ⁷ School of Population and Public Health, Faculty of Health and Medical Sciences, University of Western Australia, Nedlands, WA 6009, Australia; jane.heyworth@uwa.edu.au
 - ⁸ Department of Basic Sciences, Faculty of Allied Health Sciences, University of Peradeniya, Peradeniya 20400, Sri Lanka
- * Correspondence: s.dharmage@unimelb.edu.au



Citation: Vakalopoulos, A.; Dharmage, S.C.; Dharmaratne, S.; Jayasinghe, P.; Lall, O.; Ambrose, I.; Weerasooriya, R.; Bui, D.S.; Yasaratne, D.; Heyworth, J.; et al. Household Air Pollution from Biomass Fuel for Cooking and Adverse Fetal Growth Outcomes in Rural Sri Lanka. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1878. <https://doi.org/10.3390/ijerph18041878>

Academic Editor: Paul B. Tchounwou
Received: 20 December 2020
Accepted: 11 February 2021
Published: 15 February 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: The aim of this study was to investigate the impact of biomass fuel for cooking on adverse fetal growth outcomes in Sri Lanka. A cross-sectional study of mothers recruited at maternity clinics in rural communities in Sri Lanka's Central Province was undertaken. Data pertaining to household air pollution and fetal growth parameters were collected using an interviewer-administered questionnaire. Logistic regression models, adjusted for potential confounders, were used to evaluate the impact of biomass fuel for cooking on low birth weight (LBW) and small for gestational age (SGA) parameters. Findings showed that exposure to biomass cooking fuels during pregnancy was associated with an increased risk of LBW adjusted odds ratio (aOR) 2.74 (95% CI 1.08–6.96) and SGA (aOR: 1.87, 95% CI 1.03–3.41) compared with the use of clean energy. The risk of LBW was highest for traditional biomass stoves compared to improved biomass stoves (aOR: 3.23, 95% 1.17–8.89) and biomass use in kitchens without a chimney compared to kitchens with a chimney (aOR: 4.63, 95% 1.54–13.93). Similar trends were observed for SGA.

Keywords: household air pollution; indoor air pollution; biomass; pregnancy; Sri Lanka; LBW; SGA

1. Introduction

Household air pollution (HAP) accounts for 3% of global deaths each year and 6% in low- and middle-income countries (LMICs) [1]. HAP is attributed primarily to the burning of solid fuels such as biomass for cooking [1]. Over three billion people rely upon solid fuels, including biomass, as their primary source of domestic energy worldwide [1]. Biomass refers to organic material derived from plants and animals which may be burnt to produce energy [2]. Compared to clean energies such as liquid petroleum gas (LPG), biomass fuels release higher concentrations of carbon monoxide (CO) and fine particulate matter (particulate matter < 2.5 µm in diameter (PM_{2.5})) into household environments [3]. The majority of biomass users reside in LMICs, where cost reduces access to clean energy alternatives [4].

In Sri Lanka, biomass-fuelled stoves are the leading contributor to HAP [5,6]. Sixty-six percent of Sri Lankan households opt for biomass, principally firewood, as their main source of cooking fuel [7]. Wood burning commonly occurs indoors with thermally inefficient, traditional mud stoves and in households with poor ventilation [8]. While the consequences of HAP are manifold, there is a growing body of evidence linking exposure with adverse antenatal outcomes such as low birth weight (LBW) and fetuses small for gestational age (SGA) [9–19]. Only two studies have been conducted on the association between HAP and LBW, and none on SGA in Sri Lanka to date [9,20]. The aim of this study was to investigate the association between HAP and adverse fetal growth outcomes considering primary and secondary fuel use, stove-type, and ventilation in two rural Sri Lankan communities.

2. Materials and Methods

2.1. Study Design

A cross-sectional study was undertaken using interviewer-administrated questionnaires to collect primary data on exposure, outcome and confounder variables. Questionnaires were written in English by the primary study investigators, then translated into Sinhalese and Tamil. Questions pertaining to HAP exposure were adjusted from the BOLD Core questionnaire [21]. The final survey instrument enabled many facets of HAP to be addressed simultaneously, such as choice of cooking fuel, stove type, and kitchen ventilation. Antenatal health was measured by questions formulated from birth cards including birthweight and gestational age at birth. Birth cards are provided to all pregnant mothers for each of their children at Ministry of Health (MoH) clinics nationwide.

2.2. Study Population

The source population comprised women with a child under 5 who attended a vaccination clinic at either the Udunuwara or Gampola health clinics, which are located in rural Medical Officer of Health (MoH) areas in Sri Lanka's Central Province. The total population of the two MoH areas is estimated to be 174,000 inhabitants [22]. MoH areas were selected based on them having rural and communities who were generally from a low socio-economic background. The urban community sampled was located in Gampola and housed those working or living in Sri Lanka's tea and rubber plantations. This population comprised a comparatively low socio-economic class.

2.3. Recruitment and Sampling

Recruitment occurred during child vaccination programs and routine antenatal examinations in antenatal clinics in the selected MoH areas. Mothers who attended were approached to determine eligibility. Mothers were eligible if they had at least one child aged five or under. Prospective participants were provided with detailed information on the study, and requirements of participation including a copy of the Plain Language Statement. All information and documents were provided in the primary language of the study participant. Appointments for home-visits were then scheduled. A convenience sample of 395 households was achieved, with a response rate of 100%. Out of the 24 clinics in this area, the main three clinics, which the majority of the mothers attended for vaccination, were selected to obtain a representative sample.

2.4. Data Collection

Data collection took place over five weeks between August and September 2019. Questionnaires were administered by pre-intern medical students from the University of Peradeniya. Questions focused on household fuel use, stove type, ventilation, and sources of HAP other than cooking (smoking, mosquito coil, incense burning and vaporisers). Questions were repeated for each pregnancy for mothers who had more than one child in the past six years. The recall period of six years was acceptable given the stability of cooking practices in the region [2]. Data on socio-demographic characteristics, including

household income and mother's education, were collected. Informed consent was obtained from each mother. Ethical approval was granted by the Faculty of Allied Health Sciences at the University of Peradeniya (AHS/ERC/2019/021, 30 July 2019).

2.5. Exposure Assessment

Three different measures were used to assess exposure to HAP during pregnancy. The first method compared the primary sources of cooking fuel, "biomass" and "clean energy—mainly LPG". The second method compared stove type: "traditional biomass stove" and "improved biomass stove" to "clean energy stove". The third method categorised HAP according to primary and secondary fuel type and ventilation. Many households used more than one type of fuel for cooking. Primary fuel referred to the fuel type most frequently employed. Women utilising biomass as their primary fuel in kitchens without a chimney were categorised as "very high" exposure; those using biomass as their primary fuel in kitchens with a chimney were categorised as "high" exposure; and "moderate" exposure was specific to households using biomass as a secondary source to clean energy. Those using only clean energy formed the "low" exposure reference category. Clean energy refers to electricity, biogas, or most commonly LPG.

2.6. Outcome Assessment

Antenatal outcome data were obtained from birth cards. In Sri Lanka, birth cards are provided to all mothers at the time of pregnancy, and include information pertaining to key maternal and child health indices. The selected study outcomes were specific to two indicators of fetal growth: LBW and SGA. Birth cards provided information on the birth weight of the child in kilograms, recorded at the hospital by a nurse or midwife immediately after birth. Children were then categorised as being of LBW if they were born at a weight less than 2.50 kg. Children were categorised as being SGA if they were born at a weight below the 10th percentile for a particular gestational age. SGA was determined using the World Health Organization's (WHO) internationally applicable fetal growth calculator [23]. Surveyors were able to review the birth cards of 100% of pregnancies.

2.7. Statistical Analysis

Analysis was undertaken using the STATA 15.1 (Stata Corporation, College Station, TX, USA) statistical program for both descriptive and inferential statistics. Descriptive statistics were used to summarise socio-demographic and other baseline characteristics. Logistic regression models were then used to measure the impact of HAP on LBW and SGA. Models were adjusted for confounders and clustered with respect to household to account for mothers with multiple children aged 5 or under. The most relevant confounders to the Sri Lankan context were identified using a directed acyclic graph (DAG) (Supplementary Figure S1) and are supported by the current literature [2,8,9,20,24]. These were household monthly income, education, area, the use of incense, the use of a vaporiser for mosquitos, and exposure to second-hand tobacco smoke. Consistent with published literature, ventilation was included in the models for exposure assessment methods 1 and 2 described previously, and was measured by the presence or absence of a chimney [25].

3. Results

Data were collected for 245 rural households in Udunuwara, 65 households in Gampola, and 74 households in the urban sector in Kandy district in Sri Lanka. Incomplete questionnaire responses meant limited or no data on childhood outcomes and resulted in the exclusion of five households from the analysis. A further six households were excluded due to multiple births: five cases of twins and one case of triplets. The final dataset included information on 445 live births in the past five years, 385 mothers from 384 households.

3.1. Descriptive Statistics

3.1.1. Demographics

The rural areas of Udunuwara and Gampola were home to 81% of study participants, with 45% in households using biomass fuel. The remaining 19% of participants lived in urban communities, where the prevalence of biomass use was 64%. The primary profession for three out of four mothers was housewife. For households earning over LKR 75,000 (USD 414) each month, 75% opted for non-polluting gas or electric stoves. This is compared to 39% for households with monthly earnings of under LKR 25,000 (USD 137) (Table 1).

Table 1. Household demographics according to the primary source of cooking fuel used during pregnancy by 385 mothers in Central Sri Lanka.

	Clean Energy (Unexposed) <i>n</i> = 199 (51.69%)	Biomass (Exposed) <i>n</i> = 186 (48.31%)	Total <i>n</i> = 385, (%)
Area, <i>n</i> (%)			
Rural	172 (55.31)	139 (44.69)	311 (80.78)
Urban	27 (36.49)	47 (63.51)	74 (19.22)
Education level of mother, <i>n</i> (%)			
Primary (Grades 1–6)	3 (50.00)	3 (50.00)	6 (1.56)
Secondary (Grades 7–12)	80 (41.45)	113 (58.55)	193 (50.26)
Tertiary (University level)	116 (62.70)	69 (37.30)	185 (48.18)
Occupation of mother, <i>n</i> (%)			
Housewife	145 (50.35)	143 (49.65)	288 (75.99)
Day labour	9 (37.50)	15 (62.50)	24 (6.33)
Pink collar *	15 (53.57)	13 (46.43)	28 (7.39)
White collar **	18 (78.26)	5 (21.74)	23 (6.07)
Other ***	9 (56.25)	7 (43.75)	16 (4.22)
Monthly income (LKR ****), <i>n</i> (%)			
≤25,000 (≤USD 132)	36 (39.13)	56 (60.87)	92 (24.53)
25,001–50,000 (USD 132–263)	107 (53.77)	92 (46.23)	199 (53.07)
50,001–75,000 (USD 263–395)	24 (50.00)	24 (50.00)	48 (12.80)
≥ 75,001 (≥USD 395)	27 (75.00)	9 (25.00)	36 (9.60)
Average number of household members, \bar{x}(SD)	4.87 (1.27)	5.35 (1.29)	5.11 (1.30)

* pink collar jobs include nurse, teacher, receptionist, florist, beautician, tailor; ** white collar jobs include accountant, doctor, pharmacist, officer, advisor, clerk; *** other jobs include business/shop owner, police force; **** Sri Lankan Rupee.

3.1.2. Preference for Stove and Fuel Type

The proportion of households primarily using biomass or clean energy (LPG or electricity) was 48% and 52%, respectively. All biomass-using households opted for firewood, while for 22% of these households, almost half of biomass users, supplemented firewood with the husks and the shells of coconuts. Among biomass-using households, 55% used improved mud stoves and 45% used traditional mud stoves. LPG was used in 51% of households. The majority of households (86%) employed a secondary source of cooking fuel. Similar trends in the distribution of clean energy compared to biomass was observed for secondary fuel use, although electricity demonstrated greater use, in 20% of all households (Table 2). The majority of households using clean energy (LPG or electricity) as their primary source of fuel opted for biomass as their secondary source. The majority of households using biomass as their primary source of fuel opted for clean energy (LPG or electricity) as their secondary source (Supplementary Table S3).

Table 2. Primary and secondary cooking fuels and stove types used during pregnancy by 385 mothers in Central Sri Lanka.

	Primary Use, (%) <i>n</i> = 384 (100)	Secondary Use, (%) <i>n</i> = 331 (86.18) **
Stove type		
Gas	198 (51.56)	116 (30.21)
Electric	1 (0.26)	67 (17.45)
Traditional Biomass	83 (21.62)	57 (14.84)
Improved Biomass	102 (26.56)	91 (23.70)
Fuel type *		
Biomass		
Firewood	185 (48.18)	147 (38.28)
Coconut shells/husks	84 (21.88)	55 (14.32)
Sawdust	1 (0.26)	-
Rice husks	-	1 (0.26)
Total biomass	185 (48.18)	148 (38.54)
Clean energy		
LPG	197 (51.30)	117 (30.47)
Biogas	1 (0.26)	-
Electricity	1 (0.26)	77 (20.05)
Total clean energy	199 (51.82)	183 (47.66)

* Percentages for the type of biomass fuel do not add up because some households used more than one source of fuel for the same stove type. ** Percentages in this column are specific the total number of households.

3.1.3. Cooking Practices, Kitchen Characteristics and Maternal Health Indices

Sixty percent of mothers devoted 2–3 h to cooking each day during the pregnancy period, and a further 29% cooked for more than three hours daily. Four out of five mothers engaged in cooking during all trimesters of pregnancy. Thirty percent of households that used biomass as the primary source of fuel did not have a chimney (Supplementary Table S1).

The prevalence of LBW was 13%, of which 59% were exposed to biomass emissions in utero. The prevalence of SGA was 43%, and children who were born pre-term represented 7.5% of the sample. The average birth weight was 2.92 kg, and the average gestational period was 38.8 weeks (Supplementary Table S2).

3.2. The Association between HAP and Adverse Fetal Growth

3.2.1. The Association between HAP and LBW

Mothers using biomass for cooking during pregnancy had an increased chance of delivering a child with LBW compared to those using clean energies (adjusted odds ratio (aOR) 2.74 (95% CI 1.08–6.96). Mothers using traditional biomass stoves had an increased risk of giving birth to a child with LBW compared to those using clean energy (aOR 3.23 (95% CI 1.17–8.89)). An increased risk was also seen between improved biomass stoves and LBW, but this was not statistically significant (aOR 2.11 95%CI (0.80–5.53)) (Table 3).

Very high HAP exposure (biomass as a primary fuel source in a no-chimney kitchen) resulted in an increased risk of LBW compared to the low exposure reference category (clean energy as only source of cooking fuel), with an aOR of 4.63 (95% CI 1.54–13.93) (Table 3). As exposure to HAP increased from low to very high, so did the risk of LBW (*p*-value for overall trend = 0.005) (Table 3).

Table 3. The impact of household air pollution (HAP) as determined by primary fuel type, stove type and cooking fuel, and ventilation during pregnancy on low birth weight (LBW) for 445 births in rural Sri Lanka, 2019.

Exposure to HAP	Unadjusted		Adjusted *	
	OR	95% CI	aOR	95% CI
Primary fuel *				
Biomass	1.61	0.91–2.85	2.74	1.08–6.96
Clean energy (ref)	1.00		1.00	
Primary stove type *				
Traditional Biomass	1.87	0.95–3.67	3.23	1.17–8.89
Improved Biomass	1.42	0.72–2.78	2.11	0.80–5.53
Clean stove (ref)	1.00		1.00	
Primary fuel type and kitchen ventilation**				
Very High	3.98	1.50–10.58	4.63	1.54–13.93
High	1.42	0.54–3.71	1.21	0.41–3.53
Moderate	1.41	0.54–3.73	1.19	0.41–3.43
Low (ref)	1.00		1.00	

* adjusted for income, education, area, incense, vaporiser, second-hand tobacco smoke, and chimney; ** adjusted for income, education, area, incense, vaporiser, and second-hand tobacco smoke.

3.2.2. The Association between HAP and SGA

Mothers' exposure to biomass as their primary source of cooking fuel was associated with an increased risk of giving birth to a child SGA compared to those using clean energy aOR of 1.87 (95% CI 1.03–3.41) (Table 4). Mothers' exposure to traditional biomass stoves was also associated with an increased risk of giving birth to a child SGA compared to those using clean energy stoves aOR of 2.64 (95% 1.27–4.91) (Table 4).

Table 4. The impact of HAP as determined by primary fuel type, stove type and cooking fuel, and ventilation during pregnancy on the small for gestational age (SGA) parameter for 445 births in rural Sri Lanka, 2019.

Exposure to HAP	Unadjusted		Adjusted *	
	OR	95% CI	aOR	95% CI
Primary fuel *				
Biomass	1.75	1.17–2.62	1.87	1.03–3.41
Clean energy (ref)	1.00		1.00	
Primary stove type *				
Traditional Biomass	2.39	1.41–3.80	2.64	1.27–4.91
Improved Biomass	1.39	0.86–2.25	1.26	0.78–3.06
Clean energy stove—gas or electric (ref)	1.00		1.00	
Primary fuel type and kitchen ventilation **				
Very High	2.15	1.05–4.43	1.76	0.75–4.13
High	1.68	0.93–3.07	1.60	0.80–3.21
Moderate	1.02	0.56–1.88	1.01	0.50–2.04
Low (ref)	1.00		1.00	

* adjusted for income, education, area, incense, vaporiser, second-hand tobacco smoke, and chimney; ** adjusted for income, education, area, incense, vaporiser, and second-hand tobacco smoke.

Very high exposure (biomass as a primary fuel without chimney) and high exposure (biomass as a primary fuel with chimney) was not associated with SGA. However, as exposure increased from low to very high, so did the risk of SGA. The *p*-value for this overall trend was 0.0199 (Table 4).

4. Discussion

Results from this study add to the current body of evidence that biomass fuel compared to clean energy for cooking impairs fetal growth, increasing the risk of both LBW and SGA. It is one of the few Sri-Lankan-specific studies assessing the impact of biomass on LBW and the first of its kind on SGA. Compared to other LMICs, Sri Lanka's low prevalence of competing antenatal risk factors, coupled with a high prevalence of biomass fuel makes the country an ideal study location [26]. In fact, the country achieves 100% provision of antenatal care to its citizens, and 99.5% of mothers receive skilled attendance during childbirth, standards comparable to the developed world [26,27]. Sri Lanka also has the lowest levels of ambient air pollution in South Asia [28].

Although only one other questionnaire-based investigation on the impact of wood-fuel on LBW in Sri Lanka exists to date, the results show a similar effect size to this study [20]. Comparable trends are also evident in various Indian studies [13,14] and in a meta-analysis of 19 studies, in which the use of solid fuel was associated with a 35% (95% CI: 23, 48) increased risk of LBW in LMICs [11]. In a study in Sri Lanka's Western Province, a 10-fold increase in household PM_{2.5} was associated with a 54% (aOR: 1.54, 95% CI: 1.12, 2.12) increased risk of LBW [9].

Current literature further supports findings from this study, that poor kitchen ventilation and traditional biomass stoves increase exposure to HAP, heightening the risk of adverse antenatal outcomes. According to the WHO, the target level for indoor PM_{2.5} is 25 µg/m³ [29]. A study of 53 biomass-using rural households in Sri Lanka's Western Province found 48 h averaged PM_{2.5} concentrations of 70 µg/m³ in the presence of a chimney and 371 µg/m³ without a chimney [24]. In another Sri Lanka study, personal exposure to PM_{2.5} was substantially lower for improved biomass stoves at a range of 83–114 µg/m³ compared to traditional biomass stoves at 139–215 µg/m³ [30].

4.1. Pathological Mechanisms

PM_{2.5} and CO act through various mechanisms to alter fetal health. CO binds to haemoglobin to reduce the oxygen-carrying capacity of the blood [31]. In pregnant women, this may lead to insufficient oxygen delivery to the fetus and fetal growth retardation, increasing the risk of both LBW and SGA [32]. CO is also a fetotoxin, meaning it can cross the placental barrier and act directly upon the fetus during a critical stage of development [11,32]. Physiological immaturity and elevated rates of cell proliferation makes the fetus highly sensitive to external factors [11,33].

PM_{2.5} penetrates the alveoli sacs to trigger oxidative stress and induce systemic inflammation [34]. A study of pregnant women in Nigeria found a positive association between indoor PM_{2.5} and elevated levels of the inflammatory marker TNF-α. This was reduced following a switch to clean energy [35]. A meta-analysis also found associations between prenatal exposure to PM_{2.5} and chromosomal abnormalities, notably DNA methylation of CpGs [34]. Chromosomal abnormalities have been linked to fetal growth restriction and LBW [36,37]. It is therefore possible that oxidative stress-induced epigenetic changes contribute to adverse antenatal aetiologies [38].

4.2. Strengths and Limitations

A key strength of this study was the incorporation of a wide range of HAP determinants into three different exposure models, including primary and secondary fuel use, stove type, and kitchen ventilation. In contrast, previous studies commonly limited exposure assessment to primary fuel type. Over 78% of Sri Lankan households use more than one source of cooking fuel; therefore, disregarding secondary fuel use is likely to result in residual confounding and underestimate the impact of HAP on LBW [9].

Although household questionnaires were useful in the consideration of many facets of HAP, the approach did not enable exact indoor concentrations to be measured. Using monitors to record household concentrations of PM_{2.5} and CO presents a more precise

although expensive method to measure HAP. Studies that have employed this approach drew conclusions consistent with results from this study [9,14].

Despite secondary fuel being considered in analysis, no data were collected regarding the proportionate use of secondary fuel compared to primary fuel. Such distinctions would be useful to increase the accuracy of HAP measurements using a questionnaire-based approach. Exposure data relied upon maternal recall, and although this is the most widely used method in previously published studies [9–19], it is prone to information bias. The study's participation criteria of births within the past six years aimed to limit this by reducing time of recall. Nonetheless, as a non-differential misclassification, it lends to observed associations towards the null.

Outcome data were obtained from birth cards and were therefore less susceptible to information bias. Measurements had been previously recorded by healthcare professionals who were unaware of this study. In Sri Lanka, access to public health care is widespread and 99.5% of births are facility based, prompting the attainment of reliable and inclusive data [26]. Where healthcare does not reach rural communities, selection bias would underestimate the association between HAP and LBW. This presented a limitation in previous studies conducted in other LMICs such as India and Pakistan [10,39]. A high number of home births in previous studies also resulted in a time lag between delivery and birth weight measurements, reducing data accuracy [10,13].

Another strength of this study was the consideration of SGA as an outcome of adverse fetal growth. Most prior studies limited outcome assessment to LBW. The prevalence of SGA was determined at 43%, over three times that of LBW, and is consistent with figures worldwide [40]. This difference suggests that LBW alone cannot fully account for all cases of adverse fetal growth, and that disregarding SGA would underestimate the potential burden of pregnancies impacted by HAP. Although this study has many strengths, future research may focus on objective measurements of indoor air pollutants, and other potential risk factors associated with indoor air pollution such as asthma and ear infections in children.

4.3. Policy Implications

It is no surprise that the burden of adverse fetal growth falls heavily upon the developing world. In fact, over 95% of the world's LBW children are born in LMICs, and South Asia accounts for approximately half of them [41]. HAP-induced morbidities reduce productivity and perpetuate poverty further. Women often assume household cooking activities; therefore, the disproportionate burden of HAP is also an issue of gender inequality [42]. These equity implications provide strong incentives to address the impact of biomass fuels for cooking on antenatal health worldwide.

In Sri Lanka, improvements in maternal and child health have plateaued [26]. In fact, the prevalence of LBW is 16.6% and has remained stagnant for the past 20 years [5,22]. It is clear that Sri Lanka must divert attention to other factors influencing antenatal health, such as HAP. Unfortunately, HAP is not prominent in Sri Lanka's current public health policies [43]. As evidence for the impact of HAP on birth outcomes continues to mount, there is significant opportunity to address the modifiable risk factor.

Ultimately, to mitigate the impact of exposure to biomass pollutants on fetal growth, a transition to clean energy is recommended. Unfortunately, many LMICs, including Sri Lanka, are far behind in achieving these targets due to economic, infrastructural, logistic, and cultural constraints. Therefore, as economic barriers mean a switch to clean energy is not always possible, results from this study show that improved mud stoves and increased kitchen ventilation present an alternate avenue to reduce HAP exposure until widespread access to clean energy is possible. Exposure may be further mitigated through community-based programs that aim to increase awareness of HAP and stimulate behavioural changes.

5. Conclusions

Results from this study show that exposure to HAP from the use of biomass fuel during pregnancy is associated with an increased risk of LBW and SGA compared to the use of clean energies. Traditional biomass stoves and kitchens without a chimney increased this risk further.

To improve antenatal health prospects, findings should stimulate policy and behavioural change for clean energy adoption amongst communities and government organizations. With a high prevalence of biomass use in Sri Lanka and across the developing world, results from this study have implications on both domestic and global scales.

Supplementary Materials: The following are available online at <https://www.mdpi.com/1660-4601/18/4/1878/s1>, Figure S1: Directed Acyclic Graph of confounders for the relationship between Hap and impaired fetal growth, Table S1: Household characteristics and maternal cooking habits during pregnancy for 445 live births in Central Sri Lanka in the past 6 years, Table S2: Maternal and child health indices according to the primary fuel used during pregnancy for 445 live births in Central Sri Lanka in the past 6 years, Table S3: Secondary fuels used based on primary fuel type.

Author Contributions: Conceptualization, A.V., S.C.D., J.H., S.D. and G.B.; methodology, A.V., S.C.D., J.H., S.D., O.L., P.J., I.A. and G.B.; formal analysis, A.V., O.L. and I.A.; investigation, A.V., S.C.D., J.H., S.D., R.W., D.Y., D.S.B. and G.B.; writing—review and editing, A.V., S.C.D., J.H., S.D., R.W., D.Y., P.J., D.S.B. and G.B.; visualization, A.V.; supervision, S.C.D., J.H., S.D. and G.B.; project administration, G.B., A.V., O.L., P.J. and I.A.; funding acquisition, J.H., S.C.D., S.D. and G.B. All authors have read and agreed to the published version of the manuscript.

Funding: The research was funded by the Australian Federal Government New Colombo Plan Scheme.

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Ethics Review Committee of Faculty of Allied Health Sciences, University of Peradeniya, Sri Lanka ((AHS/ERC/2019/021, 30 July 2019).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available in supplementary material.

Acknowledgments: We would like to thank the University of Peradeniya and the National Institute of Fundamental Studies in Sri Lanka for supporting this research project. Notable thanks to the university's pre-intern medical students for their involvement in data collection. Thank you also to Arjuna Thilakaratne, Provincial Director of Health Services, Central Province; MoH Udunuwara; and MoH Gampola for allowing us to conduct research in the area, and to the Sri Lankan communities who welcomed us into their homes and took time to participate in the project. A further thanks to Mahesh Senaratna and Sajith Priyankara for statistical support. A massive thanks to the Australian Federal Government and the New Colombo Plan Scholarship Scheme, who provided the funds to make the overseas aspect of this project possible. To the University of Melbourne, namely the Allergy and Lung Health Unit at the Melbourne School of Population and Global Health, thank you for providing a supportive learning environment in which students are exposed to new ideas and encouraged with valuable feedback for their own projects.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Landrigan, P.J.; Fuller, R.; Acosta, N.J.; Adeyi, O.; Arnold, R.; Baldé, A.B.; Bertollini, R.; Bose-O'Reilly, S.; Boufford, J.I.; Breyse, P.N.; et al. The Lancet Commission on pollution and health. *Lancet* **2018**, *391*, 462–512. [[CrossRef](#)]
2. Nandasena, S.; Wickremasinghe, A.R.; Sathiakumar, N. Biomass Fuel Use for Cooking in Sri Lanka: Analysis of Data from National Demographic Health Surveys. *Am. J. Ind. Med.* **2012**, *55*, 1122–1128. [[CrossRef](#)]
3. Kim, K.H.; Jahan, S.A.; Kabir, E. A review of diseases associated with household air pollution due to the use of biomass fuels. *J. Hazard. Mater.* **2011**, *192*, 425–431. [[CrossRef](#)]
4. Ritchie, H.; Roser, M. Indoor Air Pollution. Our World in Data. 2013. Available online: <https://ourworldindata.org/indoor-air-pollution> (accessed on 24 June 2020).
5. Adair-Rohani, H.; Lewis, J. *Burning Opportunity: Clean Household Energy for Health, Sustainable Development, and Wellbeing of Women and Children*; World Health Organization: Geneva, Switzerland, 2016.

6. Nandasena, Y.L.S.; Wickremasinghe, A.R.; Sathiakumar, N. Air pollution and health in Sri Lanka: A review of epidemiologic studies. *BMC Public Health* **2010**, *10*, 300. [CrossRef] [PubMed]
7. Department of Census and Statistics (DSC). *Household Population and Housing Characteristics; Census of Population and Housing*; Department of Census and Statistics (DSC): Battaramulla, Sri Lanka, 2017.
8. Elledge, M.; Nandasena, S.; Phillips, M.; Thornburg, V. *Environmental Health Risk and the Use of Biomass Stoves in Sri Lanka*; RTI Press: Research Triangle Park, NC, USA, 2010. [CrossRef]
9. Tipre, M.; Wickremasinghe, R.; Nandasena, S.; Kasturiratne, A.; Larson, R.; Meleth, S.; Wimalasiri, U.; Smith, T.; Lungu, C.; Sathiakumar, N. Prenatal exposure to household air pollution and adverse birth outcomes among newborns in Sri Lanka. *bioRxiv* **2018**, 461632. [CrossRef]
10. Wylie, B.J.; Coull, B.A.; Hamer, D.H.; Singh, M.P.; Jack, D.; Yeboah-Antwi, K.; Sabin, L.; Singh, N.; MacLeod, W.B. Impact of biomass fuels on pregnancy outcomes in central East India. *Environ. Health* **2014**, *13*, 1. [CrossRef]
11. Amegah, A.K.; Quansah, R.; Jaakkola, J.J.K. Household Air Pollution from Solid Fuel Use and Risk of Adverse Pregnancy Outcomes: A Systematic Review and Meta-Analysis of the Empirical Evidence. *PLoS ONE* **2014**, *12*, e0113920. [CrossRef]
12. Abusalah, A.; Gavana, M.; Haidich, A.B.; Smyrnakis, E.; Papadakis, N.; Papanikolaou, A.; Benos, A. Low birth weight and prenatal exposure to indoor pollution from tobacco smoke and wood fuel smoke: A matched case-control study in Gaza Strip. *Matern. Child Health J.* **2012**, *16*, 1718–1727. [CrossRef]
13. Tielsch, J.M.; Katz, J.; Thulasiraj, R.D.; Coles, C.L.; Sheeladevi, S.; Yanik, E.L.; Rahmathullah, L. Exposure to indoor biomass fuel and tobacco smoke and risk of adverse reproductive outcomes, mortality, respiratory morbidity and growth among newborn infants in south India. *Int. J. Epidemiol.* **2009**, *38*, 1351–1363. [CrossRef] [PubMed]
14. Balakrishnan, K.; Ghosh, S.; Thangavel, G.; Sambandam, S.; Mukhopadhyay, K.; Puttaswamy, N.; Sadasivam, A.; Ramaswamy, P.; Johnson, P.; Kuppuswamy, R.; et al. Exposures to fine particulate matter (PM_{2.5}) and birthweight in a rural-urban, mother-child cohort in Tamil Nadu, India. *Environ. Res.* **2018**, *161*, 524–531. [CrossRef] [PubMed]
15. Amegah, A.K.; Jaakkola, J.J.; Quansah, R.; Norgbe, G.K.; Dzodzomenyo, M. Cooking fuel choices and garbage burning practices as determinants of birth weight: A cross-sectional study in Accra, Ghana. *Environ. Health* **2012**, *11*, 78. [CrossRef] [PubMed]
16. Epstein, M.B.; Bates, M.N.; Arora, N.K.; Balakrishnan, K.; Jack, D.W.; Smith, K.R. Household fuels, low birth weight, and neonatal death in India: The separate impacts of biomass, kerosene, and coal. *Int. J. Hyg. Environ. Health* **2013**, *216*, 523–532. [CrossRef]
17. Thompson, L.M.; Bruce, N.; Eskenazi, B.; Diaz, A.; Pope, D.; Smith, K.R. Impact of reduced maternal exposures to wood smoke from an introduced chimney stove on newborn birth weight in rural Guatemala. *Environ. Health Perspect.* **2011**, *119*, 1489–1494. [CrossRef]
18. Alexander, D.A.; Northcross, A.; Karrison, T.; Morhasson-Bello, O.; Wilson, N.; Atalabi, O.M.; Dutta, A.; Adu, D.; Ibigbami, T.; Olamijulo, J.; et al. Pregnancy outcomes and ethanol cook stove intervention: A randomized-controlled trial in Ibadan, Nigeria. *Environ. Int.* **2018**, *111*, 152–163. [CrossRef] [PubMed]
19. Pope, D.P.; Mishra, V.; Thompson, L.; Siddiqui, A.R.; Rehfuess, E.A.; Weber, M.; Bruce, N.G. Risk of Low Birth Weight and Stillbirth Associated With Indoor Air Pollution From Solid Fuel Use in Developing Countries. *Epidemiol. Rev.* **2010**, *32*, 70–81. [CrossRef]
20. Pathirathna, M.L.; Abeywickrama, H.M.; Sekijima, K.; Sadakata, M.; Fujiwara, N.; Muramatsu, Y.; Wimalasiri, K.; Jayawardene, U.; De Silva, D.; Dematawewa, C. Effects of Prenatal Tobacco and Wood-Fuel Smoke Exposure on Birth Weight in Sri Lanka. *Healthcare* **2017**, *5*, 64. [CrossRef]
21. BOLD (Burden of Obstructive Lung Disease). *Core Questionnaire*; Imperial College London: London, UK, 2006.
22. Department of Census & Statistics. *Demographic and Health Survey Report: Infant and Child Mortality*; Department of Census & Statistics: Battaramulla, Sri Lanka, 2016.
23. Fetal Growth Calculator. Available online: <http://srhr.org/fetalgrowthcalculator/#/> (accessed on 6 December 2019).
24. Chartier, R.; Phillips, M.; Mosquin, P.; Elledge, M.; Bronstein, K.; Nandasena, S.; Thornburg, V.; Thornburg, J.; Rodes, C. A comparative study of human exposures to household air pollution from commonly used cookstoves in Sri Lanka. *Indoor Air* **2017**, *27*, 147–159. [CrossRef]
25. Ranathunga, N.; Perera, P.; Nandasena, S.; Sathiakumar, N.; Kasturiratne, A.; Wickremasinghe, R. Effect of household air pollution due to solid fuel combustion on childhood respiratory diseases in a semi urban population in Sri Lanka. *BMC Pediatr.* **2019**, *19*, 306. [CrossRef] [PubMed]
26. World Health Organization. *Paradox of Healthcare in Sri Lanka: A Snapshot of the Last Decade from a Partnership of Sixty Years*; World Health Organization: Colombo, Sri Lanka, 2014.
27. Siriwardhana, D.D.; Pathmeswaran, A.; Wickremasinghe, A.R. Socioeconomic inequality and determinants of postnatal home visits made by public health midwives: An analysis of the Sri Lanka Demographic and Health Survey. *PLoS ONE* **2019**, *14*, e0215816. [CrossRef]
28. Krishna, B. Tackling the Health Burden of Air Pollution in South Asia. *BMJ Br. Med. J.* **2017**, 359. Available online: <https://www.jstor.org/stable/26950677> (accessed on 19 January 2021). [CrossRef]
29. WHO. WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide. 2006. Available online: https://apps.who.int/iris/bitstream/handle/10665/69477/WHO_SDE_PHE_OEH_06.02_eng.pdf (accessed on 13 March 2019).

30. Phillips, M.J.; Smith, E.A.; Mosquin, P.L.; Chartier, R.; Nandasena, S.; Bronstein, K.; Elledge, M.F.; Thornburg, V.; Thornburg, J.; Brown, L.M. Sri Lanka Pilot Study to Examine Respiratory Health Effects and Personal PM_{2.5} Exposures from Cooking Indoors. *Int. J. Environ. Res. Public Health* **2016**, *13*, 791. [[CrossRef](#)]
31. Vrijens, K.; Tsamou, M.; Madhloum, N.; Gyselaers, W.; Nawrot, T.S. Placental hypoxia-regulating network in relation to birth weight and ponderal index: The ENVIRONAGE Birth Cohort Study. *J. Transl. Med.* **2018**, *16*, 2. [[CrossRef](#)]
32. WHO. *Indoor Air Pollution from Solid Fuels and Risk of Low Birth Weight and Stillbirth: Report from a Symposium Held at the Annual Conference of the International Society for Environmental Epidemiology (ISEE)*; World Health Organization: Johannesburg, South Africa, 2005.
33. Nieuwenhuijsen, M.J.; Dadvand, P.; Grellier, J.; Martinez, D.; Vrijheid, M. Environmental risk factors of pregnancy outcomes: A summary of recent meta-analyses of epidemiological studies. *Environ. Health* **2013**, *12*, 6. [[CrossRef](#)]
34. Gruzieva, O.; Xu, C.J.; Yousefi, P.; Relton, C.; Merid, S.K.; Breton, C.V.; Gao, L.; Volk, H.E.; Feinberg, J.I.; Ladd-Acosta, C.; et al. Prenatal Particulate Air Pollution and DNA Methylation in Newborns: An Epigenome-Wide Meta-Analysis. *Environ. Health Perspect.* **2019**, *127*, 057012. [[CrossRef](#)] [[PubMed](#)]
35. Olopade, C.O.; Frank, E.; Bartlett, E.; Alexander, D.; Dutta, A.; Ibigbami, T.; Adu, D.; Olamijulo, J.; Arinola, G.; Karrison, T.; et al. Effect of a clean stove intervention on inflammatory biomarkers in pregnant women in Ibadan, Nigeria: A randomized controlled study. *Environ. Int.* **2017**, *98*, 181–190. [[CrossRef](#)] [[PubMed](#)]
36. Quenby, S. Recurrent miscarriage. *Obstet. Gynaecol. Reprod. Med.* **2010**, *20*, 306–310. [[CrossRef](#)]
37. Boghossian, N.S.; Horbar, J.D.; Carpenter, J.H.; Murray, J.C.; Bell, E.F. "Major Chromosomal Anomalies among Very Low Birth Weight Infants in the Vermont Oxford Network. *J. Pediatr.* **2012**, *160*, 774–780.e11. [[CrossRef](#)] [[PubMed](#)]
38. Backes, C.H.; Nelin, T.; Gorr, M.W.; Wold, L.E. Early Life Exposure to Air Pollution: How Bad Is It? *Toxicol. Lett.* **2013**, *216*, 47–53. [[CrossRef](#)]
39. Siddiqui, A.R.; Peerson, J.; Brown, K.H.; Gold, E.B.; Lee, K.; Bhuta, Z.A. Indoor air pollution from solid fuel use and low birth weight (LBW) IN PAKISTAN. *Epidemiology* **2005**, *16*, S86. [[CrossRef](#)]
40. Black, R.E. Global Prevalence of Small for Gestational Age Births. In *Low-Birthweight Baby: Born Too Soon or Too Small*; Karger Publishers: Berlin, Germany, 2015; Volume 81, pp. 1–7. [[CrossRef](#)]
41. Cutland, C.L.; Lackritz, E.M.; Mallett-Moore, T.; Bardaji, A.; Chandrasekaran, R.; Lahariya, C.; Nisar, M.I.; Tapia, M.D.; Pathirana, J.; Kochhar, S.; et al. Low birth weight: Case definition & guidelines for data collection, analysis, and presentation of maternal immunization safety data. *Vaccine* **2017**, *35*, 6492–6500. [[CrossRef](#)]
42. Desai, M.A.; Mehta, S.; Smith, K.R. *Indoor Smoke from Solid Fuels: Assessing the Environmental Burden of Disease at National and Local Levels*; World Health Organization: Geneva, Switzerland, 2004.
43. Elledge, M.F.; Phillips, M.J.; Thornburg, V.E.; Everett, K.H.; Nandasena, S. A profile of biomass stove use in Sri Lanka. *Int. J. Environ. Res. Public Health* **2012**, *9*, 1097–1110. [[CrossRef](#)]