











# Fungal Bioluminescence: Past, Present, and Future

Wenhua Lu <sup>1,2</sup> , Alviti Kankanamalage Hasith Priyashantha <sup>2</sup> , Mahesh C. A. Galappaththi <sup>3</sup> ,  
Saowaluck Tibpromma <sup>1</sup> , Dong-Qin Dai <sup>1</sup> , Nimesha M. Patabendige <sup>4</sup> , Bhagya M. Premarathne <sup>5</sup>,  
Dinesh M. Kulasuriya <sup>6</sup>, Aseni N. Ediriweera <sup>7</sup>, Thilina S. Nimalrathna <sup>8,9,10,11</sup>, Nakarin Suwannarach <sup>2,12</sup> ,  
Saisamorn Lumyong <sup>2,12</sup> , Alvin Tang <sup>13</sup> , Shi-Cheng Shao <sup>8,\*</sup> and Samantha C. Karunarathna <sup>1,\*</sup> 

- <sup>1</sup> Center for Yunnan Plateau Biological Resources Protection and Utilization, College of Biological Resource and Food Engineering, Qujing Normal University, Qujing 655011, China; reuven0319@gmail.com (W.L.); saowaluckfai@gmail.com (S.T.); cicaidongqin@gmail.com (D.-Q.D.)
- <sup>2</sup> Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand; priyashanthahasith@gmail.com (A.K.H.P.); suwan.462@gmail.com (N.S.); scboi009@gmail.com (S.L.)
- <sup>3</sup> Harry Butler Institute, Murdoch University, Perth, WA 6150, Australia; mcagalappaththi@gmail.com
- <sup>4</sup> School of Medical, Molecular and Forensic Sciences, Murdoch University, Murdoch, WA 6150, Australia; nimesha.patabendige@gmail.com
- <sup>5</sup> National Institute of Fundamental Studies (NIFS), Hanthana, Kandy 20000, Sri Lanka; bmadhushani755@gmail.com
- <sup>6</sup> Postgraduate Institute of Science (PGIS), University of Peradeniya, Peradeniya 20400, Sri Lanka; dinesh1madusanka@gmail.com
- <sup>7</sup> Centre for Mountain Futures, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China; aseniediriweera@gmail.com
- <sup>8</sup> CAS Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Xishuangbanna 666303, China; tnimalrathna@gmail.com
- <sup>9</sup> Southeast Asia Biodiversity Research Institute, Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Xishuangbanna 666303, China
- <sup>10</sup> Yunnan International Joint Laboratory of Southeast Asia Biodiversity Conservation, Yunnan Key Laboratory for Conservation of Tropical Rainforests and Asian Elephants, Xishuangbanna 666303, China
- <sup>11</sup> University of Chinese Academy of Sciences, Beijing 100101, China
- <sup>12</sup> Center of Excellence in Microbial Diversity and Sustainable Utilization, Chiang Mai University, Chiang Mai 50200, Thailand
- <sup>13</sup> Continuing and Professional Education Division, School of Continuing Education, The Hong Kong Baptist University, Hong Kong 999077, China; alvintang@hkbu.edu.hk
- \* Correspondence: shaoshicheng@xtbg.org.cn (S.-C.S.); samanthakarunarathna@gmail.com (S.C.K.)



**Citation:** Lu, W.; Priyashantha, A.K.H.; Galappaththi, M.C.A.; Tibpromma, S.; Dai, D.-Q.; Patabendige, N.M.; Premarathne, B.M.; Kulasuriya, D.M.; Ediriweera, A.N.; Nimalrathna, T.S.; et al. Fungal Bioluminescence: Past, Present, and Future. *Diversity* **2024**, *16*, 539.  
<https://doi.org/10.3390/d16090539>

Academic Editor: Michael Wink

Received: 25 July 2024

Revised: 25 August 2024

Accepted: 27 August 2024

Published: 2 September 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

**Abstract:** The complex and diverse phenomenon of fungal bioluminescence has captured human curiosity. Nevertheless, in the field of studies, there are not many attempts made particularly to reveal the new species of these interesting fungi. This study comprehensively reviews the diversity, distribution, evolution, bioluminescence mechanisms, ecological roles, and potential applications of these fungi. Most importantly, we also present an updated list of the reported bioluminescent fungi (122) so far identified from five distinct evolutionary lineages worldwide—*Armillaria*, *Eoscyphella*, *Lucentipes*, *Mycenoid*, and *Omphalotus*—mainly in tropical and subtropical areas. Bioluminescent fungi are descended from the last common ancestor of the *Mycenoid* and the *Marasmioid* clades of *Agaricales*, which have been maintained for at least 160 million years of evolution. We underscore the potential for future research to understand the ecological role of bioluminescent fungi, inspiring hope and optimism for the future of this field.

**Keywords:** biodiversity; bioluminescence; distribution; luciferin-luciferase; mushrooms

## 1. Introduction

### 1.1. Bioluminescence as a Common Term

Bioluminescence, a captivating natural phenomenon, is found in various living organisms. ‘Bio’ (in Greek) stands for life, while ‘lumen’ (in Latin) means light. Bioluminescence

is the emission of light from living organisms, a type of chemiluminescence that produces light without heat due to chemical reactions (luciferase-catalyzed oxidation reaction of luciferin) accompanied by energy stores, enzymes, substrates, and other molecules [1,2]. It is thought to represent the distribution of bioluminescent organisms throughout the Tree of Life across approximately 17 phyla and over 700 genera [3,4].

Further, bioluminescence manifests independently in at least 94 origins in the Tree of Life [5]. Throughout it, researchers have identified 40 distinct bioluminescent systems, with 13 pairs of luciferin–luciferase being the most thoroughly studied, understood, and practically applied [2,6,7]. However, the full extent of bioluminescent diversity is yet to be uncovered.

### 1.2. Brief History of Studies on Bioluminescent Organisms

Looking back at the historical records of bioluminescence, Harvey's early documents traced bioluminescent organisms back to ancient Greece and Rome [8,9]. Aristotle (384–322 BCE) stood out as a pioneer, making groundbreaking observations and recognizing the self-luminosity of these organisms. His detailed records included observations on dead fish, bioluminescent bacteria in the flesh of fish, and the bioluminescence of fireflies and worms [10,11].

Pliny the Elder's *Naturalis Historia* (23–79 CE) provided the first specific and comprehensive record of bioluminescent organisms [9]. Despite lacking independent verification, beliefs in the existence of bioluminescent birds persisted for over a thousand years [10,11]. The Dark Ages in Europe (500 CE) witnessed a scientific slowdown, but literature described ocean 'phosphorescence' and mentioned the Chinese 'candle fly' [12].

The Renaissance period witnessed a significant revival of learning, with reports of the 'burning sea' and mysterious lights at sea by Christopher Columbus. Oviedo (1478–1557) documented bioluminescent organisms and Sir Francis Drake (1540–1596) observed tropical fireflies, marking a significant advancement in bioluminescence studies [10]. In the 16th century, Conrad Gessner focused studies on luminous animals, plants, and stones, further contributing to our understanding of bioluminescence [9,11].

Three landmark bioluminescence studies were conducted during the scientific revolution in the 17th century. Philosophers like Francis Bacon (1561–1626) and René Descartes laid bioluminescence foundational principles, leading to a surge of interest in luminescent phenomena. The study of Robert Boyle (1627–1691) emphasized the importance of interrogating nature through experimentation [13].

Expanding research in the field perspective globally, evidence of bioluminescent species in Eastern countries, particularly China and Japan, became apparent. Records from the Chinese Tang and Liang dynasties trace bioluminescence back centuries [12]. The first appearance of bioluminescent fungi in Japanese literature was in ancient tales in Japan's Heian Period (6–12th century) [14]. The 19th century brought a renewed focus, with significant contributions by Dubois and Harvey, shedding light on the mechanisms of bioluminescence. Since the 20th century, more precise identifications and research have been carried out thanks to methodological and technological advancements. The blooming of molecular approaches over the past 20 years has brought significant changes in fungal taxonomy, and several databases have been launched to the public, e.g., Index Fungorum [15] and MycoBank [16]. The Amsterdam Declaration on Fungal Nomenclature "One fungus = one name" [17], ITS designated as a universal barcode for fungi, and the NCBI RefSeq Targeted Loci project for ITS has initiated [18,19], obligate registration for the valid publication of new fungal names [20]. Remarkably, the progression of molecular phylogeny has unveiled an unprecedented spectrum of fungal diversity, connecting researchers worldwide. Incorporating culture-independent techniques, notably high-throughput amplicon sequencing, has substantially escalated the enumeration of fungal operational taxonomic units. Further, throughout the last two decades, numerous innovative taxa encompassing novel divisions, classes, orders, and families have been methodically established. Molecular phylogenetics, in particular, has been instrumental in morphologically similar species, thereby advanc-

ing the understanding of fungi. Correspondingly, this genomic revolution has similarly contributed to discovering and characterizing new bioluminescent mushrooms.

Today, bioluminescent fungi (e.g., *Armillaria*, *Mycena*, and *Roridomyces*) [21–23] and bacteria (e.g., *Photobacterium* and *Vibrio*) [24,25] are the most recognized microorganisms, while animals such as fishes (e.g., *Lanternfish*) [26] and insects (e.g., *Chequevaria* and *Photuris*) are also among the popularly studied groups [27,28]. Recently, the continuous identification of bioluminescent fungi has drawn enormous attention from many research groups worldwide, leading to the discovery of many novel species [29,30]. This ongoing study trend is the driving force behind this review, which aims to provide a comprehensive overview of the current state of bioluminescent research.

### 1.3. Aspects of Bioluminescent Fungi


Bioluminescent fungi, also known as glowing fungi, can be spotted in nature by emitting a green light (delayed fluorescence), generally growing on the base of dead bamboo, tree trunks, roots, decaying wood, and fallen leaves [31]. Visible at nightfall, bioluminescence can be observed in living cultures and fruiting bodies for at least days or even a week. However, in a dense forest's darkest place, they are best observed with the naked eye at nighttime. Significant progress has been made in unraveling the mysteries surrounding these bioluminescent fungi, yet certain aspects remain unresolved. Recent taxonomic studies have shown that many works have been attempted on the taxonomy and evolution of bioluminescent mushrooms, reporting more than 40 bioluminescent mushroom species in the last decade [2,22,32–35].

This review addresses the recent surge of interest in bioluminescent fungi, particularly their diversity, worldwide distribution, evolution, glowing mechanism, and ecological significance. Furthermore, this review explores the potential applications of these fascinating organisms, offering a glimpse into the exciting future of bioluminescent research. Our study meticulously screened 35 papers to compile the species list, including reviews and original articles. We conducted an extensive literature survey across various platforms and databases, such as Scopus, National Center for Biotechnology Information (NCBI), Google Scholar, and China National Knowledge Infrastructure (CNKI), using the keywords 'bioluminescence', 'bioluminescence mushroom', and 'light fungi'. We also gathered gray literature through Google's general platform. The papers were chosen based on relevance, recent publication dates, peer-review status, and citation count, ensuring a rigorous and comprehensive literature review. Note that each scientific name was cross-checked in Index Fungorum (<http://www.indexfungorum.org/>) (accessed on 5 June 2024) and Mycobank (<https://www.mycobank.org/>) (<http://www.indexfungorum.org/>) (accessed on 5 June 2024).

## 2. Diversity and Distribution of Bioluminescent Fungi

A team of fungal experts [36] recently assessed the fungal diversity in the world using four main academic pathways, *viz.* scaling laws, fungus/plant ratios, actual versus previously known number of species, and DNA-based studies; according to them, there are likely to be 2–3 million species of fungi, with a best estimate of 2.5 million. Nevertheless, the findings of these magnificent organisms are far behind; as of 2024, around 155,000 species have only been recorded and described by taxonomists, which is comparatively lower than other particular types of fungi. For example, more than 800 genera of endophytic fungi [37] and 50,000 species of mycorrhizal fungi [38] have been recognized. Currently, over 2500 species of novel fungi are named yearly; if this continues at the current rate, it will take 750–1000 years to name the remaining unknown species [36]. According to the most recent report by Stefani et al. [39], over 125 bioluminescent fungi have been highlighted. However, this study identified the presence of 122 species (see Table 1). Different parts of fungi may glow based on the species: comparatively, 37 species (30.3%) have been reported to have both fruiting body and mycelium bioluminescence, 38 species (44.7% of known bioluminescent fruiting bodies) display undetermined mycelium bioluminescence,

36 species (29.5%) present only mycelium bioluminescence, one species shows an undetermined fruiting body bioluminescence, and 48 species (39.3%) present only fruiting body bioluminescence. Furthermore, 14 species (29.1%) have not been specified where they emit light. Figure 1 shows the global distribution of all these bioluminescent fungi. Despite the regional study bias, according to the available findings, bioluminescent fungi are mainly documented in Asia, North America, and South America.



**Figure 1.** The global distribution of bioluminescent fungi is based on the available literature; as the study was conducted, the richest fungi species were reported in Asia, followed by North and South America.

**Table 1.** The list of bioluminescent fungi reported worldwide. Bioluminescence can be produced by the entire fungus or, sometimes, only through mycelia, fruiting bodies, or spores.

	Fungal Taxa	Distribution	Glowing Part					References
			Mycelium	Fruiting Bodies	Cap	Stipe	Spores	
Armillaria Lineage	<i>Armillaria borealis</i>	Russia	+	/	/	/	/	[40]
	<i>Armillaria calvescens</i>	The USA	+	/	/	/	/	[41]
	<i>Armillaria cepistipes</i>	The USA	+	/	/	/	/	[41]
	<i>Armillaria fuscipes</i>	Malaysia	+	/	/	/	/	[42]
	<i>Armillaria gallica</i>	Europe and the USA	+	/	/	/	/	[42]
	<i>Armillaria gemina</i>	The USA	+	/	/	/	/	[42]
	<i>Armillaria mellea</i>	China, Europe, India, and the USA	+	/	/	/	/	[42–44]
	<i>Armillaria nabsnona</i>	The USA	+	/	/	/	/	[41]
	<i>Armillaria novae-zelandiae</i>	New Zealand	+	/	/	/	/	[45]
	<i>Armillaria ostoyae</i>	Europe and USA	+	/	/	/	/	[42]
	<i>Armillaria sinapina</i>	The USA	+	/	/	/	/	[41]
	<i>Desarmillaria ectypa</i>	Europe	+	+	+	/	/	[46]
Eoscyphella Lineage Lucentipes Lineage	<i>Desarmillaria tabescens</i>	Europe and the USA	+	+	/	/	/	[42]
	<i>Eoscyphella luciurceolata</i>	Brazil	?	+	?	?	?	[47]
	<i>Mycena lucentipes</i>	South America and ♣	+	+	?	+	/	[42]
	<i>Gerronema viridilucens</i>	South America	+	+	+	+	/	[42]
	<i>Cruentomyces orientalis</i>	Japan	+	+	+	+	/	[14]
	<i>Dictyopanus foliicola</i>	Japan	+	+	/	/	/	[42]
	<i>Favolaschia xtbensis</i>	China	+	+	+	+	+	[48]
	<i>Favolaschia tonkinensis</i>	China	?	+	+	+	/	[49]
	<i>Favolaschia peziziformis</i>	Japan	?	+	+	+	/	[50,51]
	<i>Filoboletus manipularis</i>	Africa, China, Sri Lanka, Thailand, and ♣	?	+	/	+	/	[52–54]
	<i>Filoboletus hanedae</i>	Japan	?	+	/	+	/	[42]
	<i>Filoboletus pallens</i>	♣	?	+	?	?	?	[42]
Mycena Lineage	<i>Filoboletus yunnanensis</i>	China	?	+	?	?	/	[52,54,55]
	<i>Gerronema glutinipes</i>	Africa and China	?	+	/	/	/	[52]
	<i>Mycena abieticola</i>	Brazil	?	+	+	+	/	[55]
	<i>Mycena aspratilis</i>	Brazil and Puerto Rico	/	+	/	+	/	[55]
	<i>Mycena asterina</i>	South America	+	+	+	/	/	[42]
	<i>Mycena cahaya</i>	♣	+	+	+	+	/	[56]
	<i>Mycena chlorophos</i>	China, Japan, the Pacific Islands, Sri Lanka, and ♣	+	+	+	+	/	[42,53]
	<i>Mycena citricolor</i>	South America and the USA	+	/	/	/	/	[42]

Table 1. Cont.

Fungal Taxa	Distribution	Glowing Part					References
		Mycelium	Fruiting Bodies	Cap	Stipe	Spores	
<i>Mycena coralliformis</i>	♫	+	/	/	/	/	[53]
<i>Mycena cristinae</i>	Brazil	+	+	/	/	/	[57]
<i>Mycena crocata</i>	Switzerland	+	/	/	/	/	[58]
<i>Mycena daisyogunensis</i>	Japan	?	+	?	?	/	[42]
<i>Mycena deeptha</i>	India	+	/	/	/	/	[59]
<i>Mycena deformis</i>	Brazil	+	/	/	/	/	[60]
<i>Mycena discobasis</i>	Africa and South America	?	+	+	+	/	[42]
<i>Mycena epipterygia</i>	Europe, the USA, and Japan	+	/	/	/	/	[42]
<i>Mycena fera</i>	South America	?	+	+	+	/	[42]
<i>Mycena flammifera</i>	Japan	+	+	+	+	/	[50,51]
<i>Mycena fulgoris</i>	Mexico	/	+	/	+	/	[61]
<i>Mycena galopus</i>	Europe, the USA, and Japan	+	/	/	/	/	[42]
<i>Mycena globulispota</i>	Brazil and Mexico	?	+	/	+	/	[60,61]
<i>Mycena gombakensis</i>	♫	+	+	+	+	/	[53]
<i>Mycena guzmanii</i>	Mexico	+	+	+	+	/	[61]
<i>Mycena haematopus</i>	China, Europe, the USA, Japan, and South America	+	+	+	/	/	[42]
<i>Mycena illuminans</i>	Japan and ♫	?	+	+	/	/	[42,53]
<i>Mycena inclinata</i>	Africa, China, Europe, and the USA	+	/	/	/	/	[42]
<i>Mycena jingyinga</i>	China	+	/	/	/	/	[34]
<i>Mycena kentingensis</i>	China	+	+	+	/	/	[61]
<i>Mycena lacrimans</i>	South America	?	+	/	+	/	[42]
<i>Mycena lamprocephala</i>	Brazil	+	+	+	+	?	[62]
<i>Mycena lazulina</i>	Japan	+	+	+	+	/	[50,51]
<i>Mycena luceata</i>	Mexico	?	+	+	/	?	[23]
<i>Mycena luciferina</i>	Mexico	?	+	+	/	?	[23]
<i>Mycena lucinieblae</i>	Mexico	+	/	/	/	?	[23]
<i>Mycena luguensis</i>	China	+	/	/	/	/	[34]
<i>Mycena lumina</i>	Mexico	+	+	+	+	/	[61]
<i>Mycena luxaeterna</i>	Brazil	+	+	/	+	/	[55]
<i>Mycena luxarboricola</i>	Brazil	?	+	+	+	/	[55]
<i>Mycena lux-coeli</i>	Japan	?	+	+	+	/	[42]
<i>Mycena luxfoliata</i>	Japan	+	/	/	/	/	[50,51]
<i>Mycena luxfoliicola</i>	Mexico	+	+	+	+	/	[61]

Table 1. Cont.

Fungal Taxa	Distribution	Glowing Part					References
		Mycelium	Fruiting Bodies	Cap	Stipe	Spores	
<i>Mycena luxmanantlanensis</i>	Mexico	+	+	+	/	?	[23]
<i>Mycena luxperpetua</i>	Puerto Rico	+	+	+	+	/	[42]
<i>Mycena maculata</i>	Africa, Europe, and the USA	+	/	/	/	/	[42]
<i>Mycena margarita</i>	Belize, Dominican Republic, Jamaica, Puerto Rico, and Brazil	/	+	+	+	/	[42,63]
<i>Mycena nebula</i>	Mexico	?	+	+	+	/	[61]
<i>Mycena nocticaelum</i>	♣	+	+	+	/	/	[53]
<i>Mycena noctilucens</i>	Pacific Islands and ♣	?	+	+	+	/	[42,53]
<i>Mycena oculisymphae</i>	Brazil	/	+	+	+	/	[60]
<i>Mycena olivaceomarginata</i>	Europe and the USA	+	/	/	/	/	[42]
<i>Mycena perlae</i>	Mexico	/	+	+	/	/	[61]
<i>Mycena polygramma</i>	China, Europe, the USA, Japan, and Africa	+	+	/	/	/	[42]
<i>Mycena pseudostylobates</i>	Japan	+	?	?	?	/	[42]
<i>Mycena pura</i>	China, Europe, the USA, Japan, and South America	+	/	/	/	/	[42]
<i>Mycena rosea</i>	Europe	+	/	/	/	/	[42]
<i>Mycena roseoflava</i>	New Zealand	+	+	/	+	/	[45]
<i>Mycena sanguinolenta</i>	China, Europe, the USA, and Japan	+	/	/	/	/	[42]
<i>Mycena seminau</i>	♣	+	+	+	/	/	[56]
<i>Mycena silvaelucens</i>	♣	?	+	+	+	/	[42,56]
<i>Mycena sinar</i>	♣	+	+	+	+	/	[56]
<i>Mycena singeri</i>	South America and ♣	?	+	+	+	/	[42]
<i>Mycena sophiae</i>	Mexico	+	/	/	/	?	[23]
<i>Mycena</i> sp. (PDD 80772)	New Zealand	?	+	/	/	/	[42]
<i>Mycena</i> sp. (SP #380150)	South America	+	+	/	/	/	[42,64]
<i>Mycena</i> sp. (SP #380281)	South America	?	+	/	/	/	[42,64]
<i>Mycena stellaris</i>	Japan	+	+	+	+	/	[50,51]
<i>Mycena stylobates</i>	Africa, China, Europe, the USA, and Japan	+	/	/	/	/	[42]
<i>Mycena tintinnabulum</i>	Europe	+	/	/	/	/	[42]
<i>Mycena venus</i>	China	+	/	/	/	/	[34]
<i>Mycena zephyrus</i>	Europe	+	/	/	/	/	[42]
<i>Panellus luminescens</i>	♣	+	+	+	+	?	[65,66]
<i>Panellus luxfilamentus</i>	Sri Lanka and ♣	+	/	/	/	/	[56]
<i>Panellus pusillus</i>	Africa, Australasia, China, Japan, the USA, South America, and ♣	?	+	?	?	/	[42,49]




Table 1. Cont.

Fungal Taxa	Distribution	Glowing Part					References
		Mycelium	Fruiting Bodies	Cap	Stipe	Spores	
	Africa, Australasia, China, Europe, Japan, the USA, and South America	+	+	+	/	/	[42]
	Japan	?	+	+	+	/	[50,51]
	Brazil	+	/	/	/	/	[60]
	Australasia	/	+	+	/	?	[42]
	Brazil, Ceylon, Malaysia, and Papua New Guinea, Singapore, and Trinidad	/	+	/	/	+	[67]
	India	?	+	/	+	/	[10,22]
	Australasia and ♣	+	+	+	+	?	[42,53]
	China, Europe, the USA, South America, and Japan	+	/	/	/	/	[42,68]
	Indonesia and ♣	/	+	+	+	/	[42]
	China	+	+	+	+	/	[69]
	Japan	+	+	/	/	/	[50,51]
	Japan	?	+	+	+	/	[50,51]
	South America	?	+	+	+	/	[42]
	Australasia, China, South America, Thailand, ♣, and ♣	?	+	+	+	/	[42,53,54]
	Japan	?	+	/	/	/	[42]
	China and Japan	+	+	/	/	/	[42,54,70]
Omphalotus Lineage	Europe and the USA	+	+	+	/	/	[42]
	China	?	+	+	/	/	[42,54,71]
	Australasia	?	+	+	+	/	[42]
	China and Europe	+	+	+	+	/	[42,54]
	The USA	/	+	/	/	/	[42]
	The USA	?	+	/	/	?	[72]
	♣	?	+	/	/	/	[42]
Ascomycota	Japan	?	+	/	/	/	[50,51]
	Europe	?	+	?	?	/	[73]

Note: For *Armillaria novae-zelandiae*, *Mycena roseoflava*, and *Omphalotus subilludens*, there is no published literature for them as bioluminescent mushrooms; however, mushroom hunters have posted glowing mushroom photos of those species. ♣ = refers to distribution within Malaysia, South Asia region; ♣ = refers to distribution within Central America and the Caribbean region; + = glow; / = do not glow; ? = no report. The original references sometimes provided only continental-level distribution information for some species, lacking specific country-level details. We have adhered to the available data for these species and recorded the distribution at the continental level. However, we have documented the specific countries accordingly for species with detailed country-level distribution information.



All bioluminescent fungi records belong to the Basidiomycota division except for *Xylaria hypoxylon* (L.) Grev., which falls under the Ascomycota [56,74]. It is worth noting that despite analyses of multiple specimens of *Xylaria hypoxylon*, differing conclusions have been drawn regarding its bioluminescence [73]; Ludwig and Gueguen reported the detection of bioluminescence in *X. hypoxylon*, whereas Molisch cultivated pure cultures for four years without observing any bioluminescent properties (for more information see [73]). This conclusion may be attributed to variations in geographical distribution and cultivation conditions. Therefore, further investigation is warranted to determine its luminescent properties conclusively [73]. Figure 2 shows five lineages that comprise all known species of bioluminescent fungi. They are part of the Mycenoid (92 species), *Armillaria* (13 species), *Omphalotus* (14 species), Lucentipes (two species), and *Eoscyphella* (one species) lineages [47].




**Figure 2.** Species percentage based on species counts reported from each lineage of bioluminescent fungi.

*Mycena* is the main genus that exhibits bioluminescence in fungi and is distributed worldwide [41,42,75]. Furthermore, accounting for the species level, for example, *Mycena chlorophos*, *M. inclinata*, and *Neonothopanus nambi* show a wide distribution worldwide [4,56,76]. Meanwhile, species like *Favolaschia xtbgensis* and *Roridomyces viridiluminus* show a restricted habitat, particularly in some places in Southwestern China (Figure 3) [48,69].

Bioluminescent fungi are not confined to a single region; their distribution spans the globe. For instance, *Gerronema viridilucens* was reported only in Brazil [77], and *Neonothopanus gardneri* from the states of Maranhão, Piauí, and Tocantins in Brazil [78]. Interestingly, some species, like *Pannellus stipticus*, are naturally found in different countries, showcasing the global nature of bioluminescent fungi. However, they did not show bioluminescence in all the recorded places. For instance, *P. stipticus* shows bioluminescence grown in North America but not in Eurasian [79]. In addition, *Armillaria mellea*, *Mycena chlorophos*, *M. deeptha*, *Nothopanus eugrammus*, *Omphalotus olearius*, *O. olivascens*, and *Roridomyces* cf. *phyllostachydis* have been reported from India [22,59,80,81]. In contrast, *Filoboletus manipularis*, *Mycena chlorophos*, and *Panellus luxfilamentus* have been found in Sri Lanka [56].

In an early study, Desjardin et al. [42] presented a revised list of bioluminescent fungi with 64 species in their distribution. Desjardin et al. [56] reported seven new luminescent fungi species two years after publication. Later, Aravindakshan et al. [59] and Shih et al. [82] reported two additional novel species from India and the Taiwan Province of China, respectively. In addition, Chew et al. [53,56] disclosed 15 bioluminescent fungi from Peninsular Malaysia, where eight were reported for the first time. Mihail [41] detected the bioluminescent mycelia of five *Armillaria* species for the first time.



**Figure 3.** Glowing piece of wood (a). Glowing mycelia on a wood in the Xishuangbanna Tropical Botanical Garden of China (b). The whole fruiting body and spores glowing in *Favolaschia xtb-gensis* in the Xishuangbanna Tropical Botanical Garden of China (c). Stipe glowing in *Roridomyces phyllostachydis* from Northeast India (d) (photo credit: Stephen Axford).

In a most recent study, Terashima et al. [50] identified another eight new species of glowing mushrooms from Japan, thus bringing the country's total reported number up to 25 [14]. Seven species of bioluminescent fungi were recorded from the cloud forests in Mexico, where six species have been identified as a new species of *Mycena*, whereas *M. globulispora* made a new distribution record for the country [61]. Furthermore, several other new species have been reported from the Taiwan Province (*M. jingyinga*, *M. luguen-sis* and *M. venus*) and Yunnan Province of China (*Favolaschia xtb-gensis* and *Roridomyces viridiluminus*) and India (*R. phyllostachydis*) [22,34,48,69]. All known bioluminescent mushrooms form gilled or poroid basidiomata within the order Agaricales. However, the latest bioluminescent species *Eoscyphella luciurceolata* represents a new lineage with a reduced cyphelloid form [47]. The discovery of new bioluminescent cyphelloid basidiomata challenges existing biological classification systems and deepens this study's understanding of bioluminescent diversity within the fungal kingdom. There is a wealth of undiscovered species, particularly in unexplored ecosystems such as forest floors, tropical regions, and polar areas, where diverse bioluminescent mushroom species may exist and represent a biodiversity hotspot for these organisms [47]. These discoveries have the potential to significantly enhance our understanding of bioluminescence mechanisms, evolutionary adaptations, and contributions to ecosystem stability.

### 3. Evolution and Mechanisms of Bioluminescent Fungi

#### 3.1. Evolution

Understanding the development of bioluminescence in fungi is a challenging and fascinating subject. Bioluminescence is most likely the result of ancient beginnings, convergent evolution, and potentially horizontal gene transfer [83]. Numerous fungal lineages have distinctive characteristics that have recently been investigated concerning the genetic and environmental factors that have influenced them [83]. Nonetheless, two main hypotheses have been proposed to explain the scattered phylogenetic distribution and lesser occurrence of bioluminescence in fungi. These hypotheses can be summarized as the

genes associated with bioluminescence shared by the common ancestor were missing in some branches and multiple convergent evolutions of bioluminescence in fungi [2]. Like other bioluminescent organisms, bioluminescent fungi have independent evolutionary occurrences, converging multiple times. Genomic analysis shows that this fragmented phylogenetic position may be another case. These findings are significant as they contribute to our understanding of the evolutionary origins and genetic mechanisms of bioluminescence in fungi, advancing mycology.

Oliveira et al. [21] uncovered a significant revelation in their research, suggesting that the origin of fungal bioluminescence can be traced back to a single evolutionary ancestry. Their evidence, demonstrated by successful light production from cross-reactions between the luciferins and luciferases of distant lineages, sheds new light on this fascinating phenomenon.

Recent studies by Kotlobay et al. [33] and Ke et al. [83] have reached a consensus, concluding that bioluminescence in fungi can be traced back to the last common ancestor of the Mycenoid and Marasmioid clades of Agaricales. This consensus echoed in recent surveys by Ke and colleagues [84], provides a solid foundation for our understanding of fungal bioluminescence.

Ke et al. [83] revisited the evolutionary dynamics of the luciferase cluster previously studied by Kotlobay et al. [33] and noted that the ancestral luciferase cluster on the same chromosome contains the genes luciferase (*Luz*), hispidin-3-hydroxylase (*H3H*), cytochrome P450 (*CYP450*), hispidin synthase (*HispS*), and caffeylpyruvate hydrolase (*CPH*). Their study corroborates and extends upon earlier findings, providing additional insights into the genes' genomic organization and evolutionary history.


Further studies revealed that gene clusters frequently undergo either deletions or retention due to differences in genomic plasticity, which explains the frequent loss of the bioluminescence property of Mycenaeen fungi [2,83,85]. The conservation of the gene cluster during the process of evolution signifies that, unlike other groups of bioluminescent organisms, bioluminescence evolves once in fungi with *Luz*, *H3H*, and *HispS* genes generated through gene duplications [33]. Moreover, the species phylogenetic tree and reconstructed phylogenetic trees of *Luz*, *H3H*, and *HispS* genes of the family Agaricaceae reveal the evolution of bioluminescent cascades in fungi. The primary *Luz* enzyme formed through a gene duplication at the base of Agaricales, followed by the duplication of *H3H* and *HispS* a few million years later [33]. These findings open up new avenues for future research, particularly in understanding the genetic mechanisms and evolutionary origins of bioluminescence in fungi.

### 3.2. Mechanisms

As previously mentioned, bioluminescent fungi, including mushrooms, have been discovered worldwide in a wide range of terrestrial environments; nevertheless, fungal bioluminescence mechanisms remain the least studied [86]. In general, bioluminescence occurs through the chemical oxidation of luciferin, catalyzed by the luciferase enzyme in the presence of oxygen [87,88]. The molecular oxygen reacts with luciferin, forming a high-energy intermediary whose decomposition emits sufficient energy to generate the emitter 'oxyluciferin' in the singlet, which is electronically excited. This excited metabolite's fluorescence property results in the emission of visible light used in nature for illumination [2,35,89].

First, it is interesting to understand how this bioluminescence mechanism has been revealed throughout history. In an early study, Dubois [90] used an in vitro luciferin/luciferase system with a mixture of heated substrate and cold enzyme–water extracts to demonstrate the first light emission experiment. In his experiment, he utilized extracts (cold and hot) from the light-emitting organs of the beetle *Pyrophorus noctilucus*. In this experimental setup, the cold extract process contained a heat-labile enzyme called luciferase that was needed to emit light. The hot extract was the thermo-stable fraction that was named luciferin. Further, it was determined that the luminescence of the mixture formed with two extracts resulted from the substrate/enzyme reaction [78].

In a study that supported Dubois [90], Airth and McElroy [91] used an in vitro setup made up of cold and hot extracts from bioluminescent fungi to confirm the role and nature of the enzymatic reaction. Later experiments by Airth and Foerster [92] explained that adding DPNH (the obsolete name for NADH) or NADPH to the cold and hot extracts activates the light emission. Furthermore, the proteinaceous cold extracts could be separated into two fractions, a pellet (insoluble) and a supernatant (soluble), by ultracentrifugation, which is necessary for the light emission in luminous fungi. Thus, the essential enzymes in each fraction for light generation postulated a two-step mechanism of enzymatic reaction [92]. Note that basidiomycetous fungi emit a green light with a maximum intensity in the 520–530 nm range [42]. Returning to Airth and Foerster's study [92], they proposed the following two-step mechanism for fungal bioluminescence:



In the first step, luciferin, a molecule that acts as an electron acceptor, is involved in the process. Here, reductase, an enzyme that catalyzes the reduction of other substances, is present in the liquid part of the mixture (supernatant). In contrast, luciferase, an enzyme that catalyzes the oxidation of luciferin, is found in the solid part (pellet). During the first step, a dark chemical reaction that does not produce a visible change occurs between NAD(P)H, a coenzyme involved in cellular respiration, luciferin, and the soluble enzyme in the supernatant. In the second step, the reduced form of luciferin reacts with molecular oxygen, catalyzed by the enzyme (luciferase) in the re-suspended pellet, producing visible light [92].

These initial findings led to understanding the chemistry behind this scenario; however, along with the development of technologies, more questions were raised, such as the in-depth aspects of the specific roles of the enzymes in bioluminescence [31,42,93]. Oliveira and Stevani [31] attempted to find the answer using an enzyme-mediated reaction by mixing a hot extract containing heat-stable substrate/luciferin with a cold extract containing the enzyme luciferase. Later, it was demonstrated that the substrates combined with the enzymes extracted from the mycelia of different bioluminescent species (*Armillaria mellea*, *Gerronema viridilucens*, *Mycena luxperpetua*, and *Neonothopanus gardneri*), and these results strongly suggest that all known bioluminescent fungi share similar types of luciferins/luciferases in bioluminescent systems [21]. Purtov et al. [94] identified the structure of fungal luciferin and its precursor as 3-hydroxyhispidin and hispidin in extracts from four diverse genera of bioluminescent fungi, namely, *Armillaria borealis*, *Mycena citricolor*, *Neonothopanus namibi*, and *Panellus stipticus*. Kaskova et al. [32] conducted an in-depth study of the mechanisms of fungal bioluminescence and color modulation and reported the structure of fungal oxyluciferin to investigate the mechanism of fungal bioluminescence. Hispidin is produced through the enzymatic activity of *HispS*. Subsequently, the resulting hispidin undergoes hydroxylation mediated by *H3H*, leading to the formation of 3-hydroxyhispidin, also known as fungal luciferin. Then, it is oxidized by  $\text{O}_2$ , generating a high-energy intermediary that decomposes in  $\text{CO}_2$  and the excited oxyluciferin. Light emission produces the ground-state oxyluciferin and hydrolyzes enzymatically into caffeic acid [32].

Kotlobay et al. [33] identified the fungal *Luz* and three other key enzymes, *HispS*, *H3H*, and *CPH*, in *Neonothopanus namibi* that jointly form the biosynthetic cycle of the fungal luciferin from caffeic acid. Fungal luciferin can be biosynthesized and recycled within this proposed mechanism. Caffeic acid is transformed to hispidin owing to *HispS* activity and is hydroxylated by *H3H*, producing 3-hydroxyhispidin fungal luciferin. The luciferase adds molecular oxygen, producing an endoperoxide (a high-energy intermediate) through decomposition that produces oxyluciferin (caffeoyl pyruvate) and light. Oxyluciferin can be recycled to caffeic acid by *CPH* [33].



In a study, Wang and Liu [95] revealed cross-reactions among four lineages of luminescent fungi, indicating that they shared a common bioluminescence mechanism, and described the bioluminescence process at the molecular level and electronic state by using multireference and density functional theory calculations. The findings revealed that fungal bioluminescence began with the cycloaddition of O<sub>2</sub> to luciferin and that formed a high-energy intermediate called  $\alpha$ -pyrone endoperoxide. This oxygenation can be explained by a charge transfer followed by a spin inversion mechanism. The high-energy intermediate thermolysis produces S1-Oxyluciferin (S1-singlet excited state) via a zwitterion intermediate. De-excitation of S1-Oxyluciferin can be a light emitter [94]. Nevertheless, according to Ke et al. [83], the complete cluster of genes involved in the bioluminescence process is still unknown to science.

In fungi, the genes that code for the enzymes that produce secondary metabolites are frequently grouped in the fungus genome [96]. In most bioluminescent fungi (e.g., *Armillaria fuscipes*, *A. mellea*, *A. ostoyae*, *A. gallica*, *Mycena citricolor*, *M. chlorophos*, *Neonothopanus nambi*, *N. gardneri*, *Omphalotus olearius*, and *Panellus stipticus*), it has been demonstrated that the aforesaid genes are generally found to be located adjacent to each other forming a cluster [33]. Rokas et al. [97] also revealed that genes in the primary and secondary metabolic pathways of fungi are often physically connected on fungal chromosomes, creating metabolic gene clusters, thus hypothesizing that this might be the reason for the formation of enzymes in the bioluminescent cascade, as this is thought to be conserved among bioluminescent fungi. In addition to the four key genes (*Luz*, *H3H*, *HispH*, and *CPH*) coding for four enzymes, *CYP450* is inside the cluster in all bioluminescent *Armillaria* and *Mycena* genomes [83]. Thus, there is controversy about the involvement of other enzymes or regulators in the bioluminescence process. On the other hand, few studies have reported the possible involvement of other genes in fungus bioluminescence [98]. In another study, Oliveira et al. [99] investigated the circadian rhythm in *Neonothopanus gardneri*. They found that the bioluminescence of the mycelium is controlled by a temperature-compensated circadian clock and the result of cycles in content/activity between the luciferase, reductase, and luciferin that comprise the bioluminescence system. Ke et al. [83] determined the regulation of bioluminescence in *Mycena kentingensis* during its development. They identified 57 gene-bearing expressions correlated to *Luz*, *H3H*, and *HispS*, agreeing with the bioluminescence mechanism discussed by Kotlobay et al. [33].

#### 4. Importance of Bioluminescent Fungi in Ecology

The ecological importance and the underlying phenomenon of bioluminescent fungi continue to be a subject of intense debate and exploration among researchers. There are several proposed hypotheses, including dispersed spores by attracting phototactic insects, deterring negative phototrophic fungivores, and potentially aposematic signals [4,100], which are a testament to the complexity of these organisms. Initially, Sivinski [101] proposed that bioluminescence in fungi serves as a warning signal to repel nocturnal fungivores or as an attractant for fungivore predators. However, Sivinski's theories challenge hypotheses suggesting that animals are primarily attracted to the aroma or odor of fungal fruiting bodies rather than their bioluminescent properties. Electroretinography is suggested to clarify whether invertebrates are attracted explicitly to bioluminescent fungi due to emitted light, although such studies are still on the bench [64].

Bioluminescent fungi have also been studied for their ability to attract insects at night for spore dispersal. Research on *Neonothopanus gardneri* has shown that the bioluminescence mechanism is regulated by circadian rhythms, involving cycles in luciferase, reductase, and luciferin activity [4,99]. This conclusion is further strengthened as researchers found that beetles bathe with the fruiting bodies of *N. gardneri* [100]. In a recent study, Karunarathna et al. [22] explained that the members of *Roridomyces* inhabiting humid environments co-evolved with some insects aiding spore dispersal. Bechara [102] demonstrated that insects such as ants, beetles, flies, wasps, and bugs are attracted to the green light emitted by bioluminescent fungi, facilitating nocturnal spore transfer in forests with min-

imal wind and high humidity. However, this hypothesis is challenged in species where bioluminescence emanates only from the stipe or mycelium [69,84]. Furthermore, some bioluminescent fungi may utilize their light as a warning signal to deter potential predators, signaling the presence of toxins or unpalatability [69,102].

Conversely, some view bioluminescence in fungi as a mere metabolic by-product devoid of ecological benefits [103,104]. Weinstein et al. [103] indicated that bioluminescence in fungi, exemplified by *Omphalotus nidiformis*, is a metabolic by-product without evident selective advantage. They suggest that the role of bioluminescence may vary among fungal lineages and environmental conditions affecting spore dispersal dynamics, such as wind patterns and insect abundance. However, the potential evolutionary advantages of bioluminescence in fungi continue to intrigue researchers, driving them to seek answers to why fungi exhibit bioluminescence. They also explore its multifaceted roles and ecological significance in different fungal lineages and environmental contexts [14,103].

## 5. Application of Fungal Bioluminescence

Fungal bioluminescence carries both historical significance and contemporary scientific potential. As aforementioned, folk stories and historical reports show that different tribes or local people, particularly in India and Indonesia, use glowing mushrooms to find their way through the dense forests [100,105,106]. In contrast, Aboriginal people in Australia consider glowing mushrooms related to the spirit [107]. In modern scientific contexts, bioluminescence has revolutionized plant biology and inspired experiments and research in biochemistry, cell biology, evolution, and photochemistry. Bioluminescence is also applied in scientific research, including several aspects such as biological sensors in environmental monitoring, effectors, hygiene control, preservation of artworks, gene assays, the detection of protein-protein interactions, bioluminescence-based imaging and photodynamic therapy, neuron treatments, and high-throughput screening in drug discovery [6,108,109].

Intriguingly, scientists are now looking for ways to switch to green light instead of light generated through electricity [100]. Recent achievements include the genetic engineering of tobacco plants (*Nicotiana tabacum* and *N. benthamiana*, see [35]) to autonomously emit light through the conversion of caffeic acid into luciferin, enabling applications in environmental assessments and potentially auto-luminescent plants [89]. The successful expression of fungal bioluminescence has also been reported in *Arabidopsis thaliana*, *Catharanthus roseus*, *Dahlia pinnata*, *Petunia hybrida*, *Rosa rubiginosa*, and *Solanum lycopersicum* [89,110,111].

Environmental bioassays can be performed using fungi's natural bioluminescent enzyme reaction [78]. Eukaryotic bioluminescent fungi are a more suitable research organism for soil toxicology than luminescent bacteria. However, the mechanism of toxicity and its specific impact on the fungal bioluminescence response is not yet fully understood. The uncoupling of oxidative phosphorylation and the depolarization of mitochondrial membranes by toxic compounds can be possible pathways; they would possibly indirectly affect the NADH availability that is involved in the bioluminescent reaction [78]. Additionally, it is still feasible to use organisms in a terrestrial setting, including bioluminescent fungi, without further engineering marker labeling [2], even if known uses include marker-labeled bacteria and marine bioluminescent *Vibrio* species [112,113]. Since 2002, toxicity tests have been developed using bioluminescent fungi, particularly with *Armillaria mellea* and *Mycena citricolor*. These studies were based on globular mycelia grown on liquid media with varying concentrations of heavy metals (copper and zinc) or chemical compounds (chlorophenol). Several bioassays with bioluminescent basidiomycetes have been developed using *Pannellus stipticus* and *Omphalotus olearius* [78]. More recently, the toxicity of the other bioluminescent lineages of *Gerronema viridilucens* and *N. gardneri* was assessed [114,115]. Bioluminescence technology plays a pivotal role in *in vivo* bioimaging, broadly applied in the study of diseases and assessing therapeutic interventions in animal models. Researchers can monitor and analyze dynamic biological processes like tumor progression and inflammatory responses in real-time by employing bioluminescent markers such as the luciferase system. This capability facilitates the evaluation of treatment efficacy

and safety profiles. The noninvasive nature, high spatiotemporal resolution, and ability to quantify drug distribution and metabolism make bioluminescence imaging indispensable for personalized medicine and innovative drug discovery [116–118]. Beyond advancing the frontiers of biomedical research, bioluminescence imaging accelerates the refinement and development of therapeutic strategies.

On the one hand, the reconstruction of the fungal bioluminescence pathway in an organism, making it autonomous luminescence, is beneficial for detecting the status of its various growth stages. It could also facilitate the development of the next generation of organic architecture, modified light-emitting plants in buildings, and urban infrastructure [33]. On the other hand, fungal bioluminescence can be indicated in agriculture when crops need water or nutrients. Due to this autonomous bioluminescence, plants can warn early about illnesses and pest attacks that could harm harvests. Furthermore, bioluminescence paves the path for eco-friendly house/street lighting, health applications, and food industries [100,105,106]. The alterations of these technologies will drive massive growth in bioluminescence in the coming future [75]. Bioluminescent mushrooms offer significant potential in both horticulture and tourism. In horticultural landscape design, they create a unique nighttime ambiance, enhancing the visual appeal of gardens through their natural glow in flowerbeds, pathways, and lawns. These mushrooms also provide educational opportunities by providing practical examples of bioluminescence in horticultural institutions. In tourism, they serve as distinctive attractions, potentially forming mushroom gardens or designated viewing areas and contributing to local economies. Incorporating bioluminescent mushrooms into eco-tourism activities, such as nocturnal ecological tours, offers visitors novel nature interactions while promoting an understanding of biodiversity and ecosystems. Overall, bioluminescent mushrooms enrich aesthetic and educational aspects and present promising opportunities for tourism development. Bioluminescent mushrooms hold tremendous promise for both horticulture and tourism. In horticultural design, these mushrooms create a magical nighttime atmosphere, adding a natural glow that enhances the beauty of gardens, whether nestled in flowerbeds, lining pathways, or dotting lawns. Beyond their aesthetic value, they also serve as practical tools in educational settings, vividly illustrating the wonders of bioluminescence in horticulture. In the realm of tourism, bioluminescent mushrooms become unique attractions, capable of forming enchanting mushroom gardens or designated viewing spots that captivate visitors and potentially bolster local economies. Integrating these mushrooms into eco-tourism initiatives, such as nocturnal ecological tours, offers guests an extraordinary opportunity to engage with nature in new and profound ways while fostering a deeper appreciation for biodiversity and ecosystems. Ultimately, bioluminescent mushrooms enrich the visual and educational dimensions and open exciting avenues for developing tourism experiences.

## 6. Conclusions and Future Directions

According to our understanding, 122 species of bioluminescent fungi have been reported. These fungi are primarily categorized within the Basidiomycota, distributed across four established phylogenetic lineages (*Armillaria*, *Lucentipes*, *Mycenoid*, and *Omphalotus*), with the recent addition of a novel fifth lineage, *Eoscyphella*. Despite phylogenetic diversity, all bioluminescent fungi share conserved bioluminescent mechanisms rooted in luciferin oxidation catalyzed by luciferase, which involves two enzymatic steps. However, detailed studies elucidating the genetic regulation of this process remain a critical area for future investigation. Recent advancements highlight the possible diverse applications of fungal bioluminescence. Bioluminescent fungi hold immense promise across a wide spectrum of disciplines, including ecology, agriculture, art, medicine, and education. They offer potential as bioindicators for environmental monitoring and innovative strategies for crop health management through genetic modification. Furthermore, their aesthetic allure has inspired creative designs such as luminous gardens and interactive art installations. In medicine, these fungi offer exciting avenues for developing new therapeutic agents and diagnostic tools. In summary, realizing the full potential of bioluminescent fungi necessi-



tates ongoing interdisciplinary collaboration. However, it is crucial to stress that continued research efforts are needed to expand the understanding of fungal biodiversity, ecological roles, and practical applications in various fields as the field constantly evolves.

**Author Contributions:** S.C.K. and S.T. conceptualized the topic; W.L. and A.K.H.P. wrote the first draft of the manuscript; M.C.A.G., N.M.P., A.N.E., B.M.P. and D.M.K. helped with the literature searches and drafted the chemistry part of the manuscript; T.S.N. prepared the map; S.T., S.-C.S., D.-Q.D., A.K.H.P., A.T., N.S., S.L. and S.C.K. edited the manuscript; S.T. and S.C.K. supervised the work; D.-Q.D., N.S., S.L., S.-C.S. and S.C.K. provided funding for the research. All authors have read and agreed to the published version of the manuscript.

**Funding:** This work was supported by the National Natural Science Foundation of China (Grant Nos. 32260004, 31760013, and 32171655), the West Light Talent Program of the Chinese Academy of Sciences (Grant No. E1XB011B01), and Chiang Mai University and Kaomai Lanna Res., Thailand (Grant No. R66IN00132). Samantha C. Karunarathna, Dong-Qin Dai, and Saowaluck Tibpromma thank the High-Level Talent Recruitment Plan of Yunnan Province (“High-End Foreign Expert” and “Young Talents” Programs) for their support.

**Data Availability Statement:** The data that support this study are available on request.

**Acknowledgments:** The authors thank the Key Laboratory of the Yunnan Provincial Department of Education of the Deep-Time Evolution on Biodiversity from the Origin of the Pearl River, for their support.

**Conflicts of Interest:** The authors declare no conflicts of interest.

## References

- Pandey, A.; Sharon, M. Bioluminescent organisms. *BAOJ Chem.* **2017**, *3*, e029.
- Ke, H.M.; Tsai, I.J. Understanding and using fungal bioluminescence—Recent progress and future perspectives. *Curr. Opin. Green Sustain. Chem.* **2022**, *33*, 100570. [[CrossRef](#)]
- Waldenmaier, H.E.; Oliveira, A.G.; Stevani, C.V. Thoughts on the diversity of convergent evolution of bioluminescence on earth. *Int. J. Astrobiol.* **2012**, *11*, 335–343. [[CrossRef](#)]
- Vieira, M.B.B.; Oliveira, I.C.; de Oliveira, M.D.D.A.; da Costa Júnior, J.S.; dos Santos, T.D.J.A.; Feitosa, C.M.; Rai, M.; Lima, N.M.; da Costa Silva, D. A review on bioluminescent fungus *Neonothopanus gardneri*. *Res. Soc. Dev.* **2022**, *11*, e16811528009. [[CrossRef](#)]
- Lau, E.S.; Oakley, T.H. Multi-level convergence of complex traits and the evolution of bioluminescence. *Biol. Rev. Camb. Philos. Soc.* **2020**, *96*, 673–691. [[CrossRef](#)] [[PubMed](#)]
- Syed, A.J.; Anderson, J.C. Applications of bioluminescence in biotechnology and beyond. *Chem. Soc. Rev.* **2021**, *50*, 5668–5705. [[CrossRef](#)] [[PubMed](#)]
- Delroisse, J.; Duchatelet, L.; Flammang, P.; Mallefet, J. Leaving the dark side? Insights into the evolution of luciferases. *Front. Mar. Sci.* **2021**, *8*, 673620. [[CrossRef](#)]
- Harvey, E.N. *Bioluminescence*; Academic Press: New York, NY, USA, 1952; pp. 1–649.
- Harvey, E.N. *A History of Luminescence: From the Earliest Times until 1900*; The American Philosophical Society: Philadelphia, PA, USA, 1957; pp. 1–692. [[CrossRef](#)]
- Lee, S.M.L. The status of bioluminescent fungal species in Singapore. *Nat. Singap.* **2022**, *1*, e2022124. [[CrossRef](#)]
- Mahish, P.K.; Chandrawanshi, N.K.; Kunjam, S.; Jadhav, S.K. Opportunities in the living lights: Special reference to bioluminescent fungi. *Energy Cris. Chall. Solut.* **2021**, *10*, 191–207.
- Lu, D. 71 species of macrofungi that bioluminescence. *Edible Med. Mushrooms* **2011**, *19*, 55–57. (In Chinese)
- Boyle, R. Observations and tryals about the resemblances and differences between a burning coal and shining wood. *Phil. Trans.* **1667**, *2*, 605–612. [[CrossRef](#)]
- Oba, Y.; Hosaka, K. The luminous fungi of Japan. *J. Fungi* **2023**, *9*, 615. [[CrossRef](#)] [[PubMed](#)]
- Kirk, P.M. World catalogue of 340K fungal names on-line. *Mycol. Res.* **2000**, *104*, 516–517.
- Crous, P.W.; Gams, W.; Stalpers, J.A.; Robert, V.; Stegehuis, G. MycoBank: An online initiative to launch mycology into the 21st century. *Stud. Mycol.* **2004**, *50*, 19–22.
- Taylor, J.W. One Fungus = One Name: DNA and fungal nomenclature twenty years after PCR. *IMA Fungus* **2011**, *2*, 113–120. [[CrossRef](#)] [[PubMed](#)]
- Schoch, C.L.; Seifert, K.A.; Huhndorf, S.; Robert, V.; Spouge, J.L.; Levesque, C.A.; Chen, W.; Fungal Barcoding Consortium; Fungal Barcoding Consortium Author List; Bolchacova, E.; et al. Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 6241–6246. [[CrossRef](#)] [[PubMed](#)]

19. Schoch, C.L.; Robbertse, B.; Robert, V.; Vu, D.; Cardinali, G.; Irinyi, L.; Meyer, W.; Nilsson, R.H.; Hughes, K.; Miller, A.N.; et al. Finding needles in haystacks: Linking scientific names, reference specimens and molecular data for Fungi. *Database* **2014**, 2014, bau061. [CrossRef] [PubMed]
20. McNeill, J.; Barrie, F.R.; Buck, W.R.; Demoulin, V.; Greuter, W.; Hawksworth, D.L.; Herendeen, P.S.; Knapp, S.; Marhold, K.; Prado, J.; et al. International Code of Nomenclature for algae, fungi, and plants. *Regnum Veg.* **2012**, 154. Available online: <http://www.iapt-taxon.org/nomen/main.php> (accessed on 25 July 2024).
21. Oliveira, A.G.; Desjardin, D.E.; Perry, B.A.; Stevani, C.V. Evidence that a single bioluminescent system is shared by all known bioluminescent fungal lineages. *Photochem. Photobiol. Sci.* **2012**, 11, 848–852. [CrossRef] [PubMed]
22. Karunarathna, S.C.; Mortimer, P.E.; Tibpromma, S.; Dutta, A.K.; Paloi, S.; Hu, Y.; Baurah, G.; Axford, S.; Marciniak, C.; Luangharn, T.; et al. *Roridomyces phyllostachydis* (Agaricales, Mycenaceae), a new bioluminescent fungus from Northeast India. *Phytotaxa* **2020**, 459, 155–167. [CrossRef]
23. Cortés-Pérez, A.; Guzmán-Dávalos, L.; Ramírez-Cruz, V.; Villalobos-Arámbula, A.R.; Ruiz-Sanchez, E.; Ramírez-Guillén, F. New Species of Bioluminescent *Mycena* Sect. *Calodontes* (Agaricales, Mycenaceae) from Mexico. *J. Fungi* **2023**, 9, 902. [CrossRef]
24. Labella, A.M.; Arahall, D.R.; Castro, D.; Lemos, M.L.; Borrego, J.J. Revisiting the genus *Photobacterium*: Taxonomy, ecology and pathogenesis. *Int. Microbiol.* **2017**, 20, 1–10.
25. Burtseva, O.; Kublanovskaya, A.; Baulina, O.; Fedorenko, T.; Lobakova, E.; Chekanov, K. The strains of bioluminescent bacteria isolated from the White Sea finfishes: Genera *Photobacterium*, *Aliivibrio*, *Vibrio*, *Shewanella*, and first luminous *Kosakonia*. *J. Photoch. Photobio. B* **2020**, 208, 111895. [CrossRef]
26. de Busserolles, F.; Marshall, N.J. Seeing in the deep-sea: Visual adaptations in lanternfishes. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **2017**, 372, 20160070. [CrossRef] [PubMed]
27. Al-Handawi, M.B.; Polavaram, S.; Kurlvskaya, A.; Commings, P.; Schramm, S.; Carrasco-López, C.; Lui, N.M.; Solntsev, K.M.; Laptinok, S.P.; Navizet, I.; et al. Spectrochemistry of firefly bioluminescence. *Chem. Rev.* **2022**, 122, 13207–13234. [CrossRef]
28. Owens, A.; Van den Broeck, M.; De Cock, R.; Lewis, S.M. Behavioral responses of bioluminescent fireflies to artificial light at night. *Front. Ecol. Evol.* **2022**, 10, 946640. [CrossRef]
29. Elkhateeb, W.A.; Daba, G.M. Bioluminescent Mushrooms: Boon for environmental health. *Environ. Sci. Arch.* **2022**, 1, 88–97. Available online: [https://www.envsciarch.com/\\_files/ugd/4b6a78\\_9155754cbd4e4412b6bab4385f8ef2e1.pdf](https://www.envsciarch.com/_files/ugd/4b6a78_9155754cbd4e4412b6bab4385f8ef2e1.pdf) (accessed on 14 October 2022).
30. Adams, C.A.; Donald, M.L.; Swearingen, C.; Escudero, E.; Sourell, S.; Landrein, S.; Seas, C.; Mueller, G.; Chaverri, P. Let there be nightlights: The ecological role of bioluminescence in a Costa Rican mushroom. *bioRxiv* **2023**. [CrossRef]
31. Oliveira, A.G.; Stevani, C.V. The enzymatic nature of fungal bioluminescence. *Photochem. Photobiol. Sci.* **2009**, 8, 1416. [CrossRef] [PubMed]
32. Kaskova, Z.M.; Dörr, F.A.; Petushkov, V.N.; Purtov, K.V.; Tsarkova, A.S.; Rodionova, N.S.; Mineev, K.S.; Guglya, E.B.; Kotlobay, A.; Baleeva, N.S.; et al. Mechanism and color modulation of fungal bioluminescence. *Sci. Adv.* **2017**, 3, e1602847. [CrossRef]
33. Kotlobay, A.A.; Sarkisyan, K.S.; Mokrushina, Y.A.; Marcet-Houben, M.; Serebrovskaya, E.O.; Markina, N.M.; Gonzalez, S.L.; Gorokhovatsky, A.Y.; Vvedensky, A.; Purtov, K.V.; et al. Genetically encodable bioluminescent system from fungi. *Proc. Natl. Acad. Sci. USA* **2018**, 115, 12728–12732. [CrossRef] [PubMed]
34. Chang, C.C.; Chen, C.Y.; Li, W.W.; Ka, W. *Mycena jingyinga*, *Mycena luguensis*, and *Mycena venus*: Three new species of bioluminescent fungi from Taiwan. *Taiwania* **2020**, 65, 396–406. Available online: <https://taiwania.ntu.edu.tw/pdf/tai.2020.65.396.pdf> (accessed on 27 July 2020).
35. Mitouchkina, T.; Mishin, A.S.; Somermeyer, L.G.; Markina, N.M.; Chepurnyh, T.V.; Guglya, E.B.; Karataeva, T.A.; Palkina, K.A.; Shakhova, E.S.; Fakhranurova, L.I.; et al. Plants with genetically encoded autoluminescence. *Nat. Biotechnol.* **2020**, 38, 944–946. [CrossRef] [PubMed]
36. Niskanen, T.; Lücking, R.; Dahlberg, A.; Gaya, E.; Suz, L.M.; Mikryukov, V.; Liimatainen, K.; Druzhinina, I.; Westrip, J.R.; Mueller, G.M.; et al. Pushing the frontiers of biodiversity research: Unveiling the global diversity, distribution, and conservation of fungi. *Annu. Rev. Environ. Resour.* **2023**, 48, 149–176. [CrossRef]
37. Rashmi, M.; Kushveer, J.S.; Sarma, V.V. A worldwide list of endophytic fungi with notes on ecology and diversity. *Mycosphere* **2019**, 10, 798–1079. [CrossRef]
38. Van Der Heijden, M.G.; Martin, F.M.; Selosse, M.A.; Sanders, I.R. Mycorrhizal ecology, and evolution: The past, the present, and the future. *New Phytol.* **2015**, 205, 1406–1423. [CrossRef] [PubMed]
39. Stevani, C.V.; Zamuner, C.K.; Bastos, E.L.; de Nóbrega, B.B.; Soares, D.M.; Oliveira, A.G.; Bechara, E.J.; Shakhova, E.S.; Sarkisyan, K.S.; Yampolsky, I.V.; et al. The living light from fungi. *J. Photochem. Photobiol. C Photochem. Rev.* **2024**, 58, 100654. [CrossRef]
40. Vydryakova, G.A.; Gusev, A.A.; Medvedeva, S.E. Effect of organic and inorganic toxic compounds on luminescence of luminous fungi. *Appl. Biochem. Microbiol.* **2011**, 47, 293–297. [CrossRef]
41. Mihail, J.D. Bioluminescence patterns among North American *Armillaria* species. *Fungal Biol.* **2015**, 119, 528–537. [CrossRef]
42. Desjardin, D.E.; Oliveira, A.G.; Stevani, C.V. Fungi bioluminescence revisited. *Photochem. Photobiol. Sci.* **2008**, 7, 170–182. [CrossRef]
43. Tan, Z.J.; Xie, D.P.; Wang, Z.; Li, W.G.; Liu, S. The study on bioluminescence condition of *Amillaria mellea*. *Acta Laser Biol. Sinica* **2001**, 10, 1007–7146. (In Chinese)

44. Patil, S.R.; Yadav, S.V. Photographic record of *Armillaria mellea* a bioluminescent fungus from Lonavala in western Ghats, India. *J. Threat. Taxa* **2022**, *14*, 20692–20694. [CrossRef]
45. BITB. Bioluminescence in the Bush: Glow in the Dark Mushrooms in Stewart Island. Available online: <https://www.myconeer.com/p/bioluminescence-in-the-bush-glow> (accessed on 24 April 2024).
46. Ainsworth, M. Searching for luminous mushrooms of the marsh fungus *Armillaria ectypa*. *Field Mycol.* **2004**, *5*, 142–144. [CrossRef]
47. Silva-Filho, A.G.; Mombert, A.; Nascimento, C.C.; Nóbrega, B.B.; Soares, D.M.; Martins, A.G.; Domingos, A.H.; Santos, I.; Della-Torre, O.H.; Perry, B.A.; et al. *Eoscyphella luciurceolata* gen. and sp. nov. (Agaricomycetes) shed light on Cyphellopsidaceae with a new lineage of bioluminescent fungi. *J. Fungi* **2023**, *9*, 1004. [CrossRef] [PubMed]
48. Nimalrathna, T.; Tibpromma, S.; Nakamura, A.; Galappaththi, M.C.A.; Xu, J.; Mortimer, P.E.; Karunarathna, S.C. The case of the missing mushroom: A novel bioluminescent species discovered within *Favolaschia* in Southwestern China. *Phytotaxa* **2022**, *539*, 244–256. [CrossRef]
49. Liu, P.G. Luminescent fungi. *Chin. Biodivers.* **1995**, *3*, 109–112. (In Chinese)
50. Terashima, Y.; Neda, H.; Hiroi, M. Luminescent intensity of cultured mycelia of eight basidiomycetous fungi from Japan. *Mushroom Sci. Biotechnol.* **2016**, *24*, 176–181.
51. Terashima, Y.; Takahashi, H.; Taneyama, Y. *The Fungal Flora in Southwestern Japan: Agarics and Boletes*; Tokai Daigaku: Kanagawa, Japan, 2016; p. 349.
52. Liu, P.G.; Yang, Z.L. Studies of classification and geographic distribution of Laschia-complex from the Southern and Southeastern Yunnan, China. *Acta Bot. Yunnanica* **1994**, *16*, 47–52. (In Chinese)
53. Chew, A.L.; Desjardin, D.E.; Tan, Y.S.; Musa, M.Y.; Sabaratnam, V. Bioluminescent fungi from peninsular Malaysia—A taxonomic and phylogenetic overview. *Fungal Divers.* **2014**, *70*, 149–187. [CrossRef]
54. Yan, J.J.; Liu, X.R.; Xie, B.G.; Deng, Y.J. Isolation, identification, and characterization of *Neonothopanus nambi* (Basidiomycota, Fungi), a new record from China. *Bull. Microbiol.* **2015**, *42*, 1703–1709.
55. Desjardin, D.E.; Perry, B.A.; Lodge, D.J.; Stevani, C.V.; Nagasawa, E. Luminescent *Mycena*: New and noteworthy species. *Mycologia* **2010**, *102*, 459–477. [CrossRef] [PubMed]
56. Chew, A.L.; Tan, Y.S.; Desjardin, D.E.; Musa, M.Y.; Sabaratnam, V. Four new bioluminescent taxa of *Mycena* sect. *Calodontes* from Peninsular Malaysia. *Mycologia* **2014**, *106*, 976–988. [CrossRef]
57. Oliveira, J.J.; Vargas-Isla, R.; Cabral, T.S.; Cardoso, J.S.; Andriolli, F.S.; Rodrigues, D.P.; Ikeda, T.; Clement, C.R.; Ishikawa, N.K. The amazonian luminescent *Mycena cristinae* sp. nov. from Brazil. *Mycoscience* **2021**, *62*, 395–405. [CrossRef]
58. Heinzelmann, R.; Baggenstos, H.; Rudolf, A. Is the bioluminescence in many *Mycena* species overlooked?—A case study from *M.crocata* in Switzerland. *Mycoscience* **2024**, *65*, MYC633. [CrossRef]
59. Aravindakshan, D.M.; Kumar, T.K.A.; Manimohan, P. A new bioluminescent species of *Mycena* sect. *Exornatae* from Kerala State, India. *Mycosphere* **2012**, *3*, 556–561. [CrossRef]
60. Desjardin, D.E.; Perry, B.A.; Stevani, C.V. New luminescent Mycenoid fungi (Basidiomycota, Agaricales) from Sao Paulo state, Brazil. *Mycologia* **2016**, *108*, 1165–1174. [CrossRef] [PubMed]
61. Cortés-Pérez, A.; Desjardin, D.E.; Perry, B.A.; Ramírez-Cruz, V.; Ramírez-Guillén, F.; Villalobos-Arámbula, A.R.; Rockefeller, A. New species, and records of bioluminescent *Mycena* from Mexico. *Mycologia* **2019**, *111*, 319–573. [CrossRef]
62. Soares, C.C.; Cabral, T.S.; Vargas-isla, R.U.B.Y.; Cardoso, J.S.; Rodrigues, D.P.; Ishikawa, N.K.; Oliveira, J.J. *Mycena lamprocephala*, a new luminescent species from the Brazilian Amazon. *Phytotaxa* **2024**, *634*, 187–203. [CrossRef]
63. Alves, M.H.; do Nascimento, C.C. *Mycena margarita* (Murrill) Murrill, 1916 (Basidiomycota: Agaricales: Mycenaceae): A bioluminescent agaric first recorded in Brazil. *Check List* **2014**, *10*, 239–243. [CrossRef]
64. Desjardin, D.E.; Capelari, M.; Stevani, C. Bioluminescent *Mycena* species from São Paulo, Brazil. *Mycologia* **2007**, *99*, 317–331. [CrossRef] [PubMed]
65. Corner, E.J.H. Descriptions of two luminous tropical agarics (*Dictyopanus* and *Mycena*). *Mycologia* **1950**, *42*, 423–431. [CrossRef]
66. Corner, E.J.H. The agaric genus *Panellus* Karst. (including *Dictyopanus* Pat.) in Malaysia. *Gard. Bull.* **1986**, *39*, 103–147. Available online: <https://biostor.org/reference/177690> (accessed on 25 July 2024).
67. Horak, E. *Mycena rorida* (Fr.) Quél. and related species from the Southern Hemisphere. *Berichte* **1978**, *88*, 20–29. [CrossRef]
68. Tolgor, B. Notes on Basidiomycetes of Jilin province (VIII). *J. Fungal Res.* **2007**, *5*, 72–74.
69. Dauner, L.A.P.; Karunarathna, S.C.; Tibpromma, S.; Xu, J.; Mortimer, P.E. Bioluminescent fungus *Roridomyces viridiluminus* sp. nov. and the first Chinese record of the genus *Roridomyces* from Southwestern China. *Phytotaxa* **2021**, *487*, 233–250. [CrossRef]
70. Yang, Z.L.; Feng, B. The genus *Omphalotus* (Omphalotaceae) in China. *Mycosystema* **2013**, *32*, 545–556.
71. Li, J.Z.; Hu, X.W. A new species of *Lampteromyces* from Hunan. *Acta Sci. Nat. Univ. Norm. Hunan* **1993**, *16*, 2.
72. OSSJ. *Omphalotus Subilludens*—Southern Jack O’ Lantern. Available online: [https://www.texasmushrooms.org/en/omphalotus\\_subilludens.htm](https://www.texasmushrooms.org/en/omphalotus_subilludens.htm) (accessed on 1 June 2024).
73. Buller, A.H.R. *Book: Researches on Fungi*; Longmans: London, UK, 1958; Volume III, pp. 416–419. Available online: <https://archive.org/details/researchesonfung03bull/page/418/mode/2up> (accessed on 30 April 2008).
74. Foerster, G.E.; Behrens, P.Q.; Airth, R.L. Bioluminescence and other characteristics of *Collybia velutipes*. *Am. J. Bot.* **1965**, *52*, 487–495. [CrossRef] [PubMed]
75. Kushwaha, V.; Hajirnis, S. A review on bioluminescent fungi: A torch of curiosity. *Int. J. Life Sci.* **2016**, *A7*, 107–110. Available online: <https://oaji.net/articles/2017/736-1518519635.pdf> (accessed on 1 December 2016).



76. Wassink, E.C. Observations on the luminescence in fungi I, including a critical review of the species mentioned as luminescent in literature. *Recl. Des Trav. Bot. Néerlandais* **1948**, *4*, 150–212.
77. Desjardin, D.E.; Capelari, M.; Stevani, C.V. A new bioluminescent Agaric from São Paulo, Brazil. *Fungal Divers.* **2005**, *18*, 9–14.
78. Stevani, C.V.; Oliveira, A.G.; Mendes, L.F.; Ventura, F.F.; Waldenmaier, H.E.; Carvalho, R.P.; Pereira, T.A. Current status of research on fungal bioluminescence: Biochemistry and prospects for ecotoxicological application. *Photochem. Photobiol.* **2013**, *89*, 1318–1326. [\[CrossRef\]](#) [\[PubMed\]](#)
79. Peterson, R.H.; Bermudes, D. Intercontinental compatibility in *Panellus stypticus* with a note on bioluminescence. *Persoonia* **1992**, *14*, 457–463.
80. Arya, C.P.; Ratheesh, S.; Pradeep, C.K. New record of luminescent *Mycena chlorophos* (Mycenaceae) from Western Ghats of India. *Stud. Fungi* **2021**, *6*, 507–513. [\[CrossRef\]](#)
81. Dutta, A.; Gupta, S.; Roy, J.K.; Ahmed, M.F. New distribution record of *Roridomyces* cf. *phyllostachydis* (Agaricales: Mycenaceae), a bioluminescent fungus from Namdapha National Park, Arunachal Pradesh, India. *J. Threat. Taxa* **2023**, *15*, 22920–22923. [\[CrossRef\]](#)
82. Shih, Y.S.; Chen, C.Y.; Lin, W.W.; Kao, H.W. *Mycena kentingensis*, a new species of luminous mushroom in Taiwan, with reference to its culture method. *Mycol. Prog.* **2014**, *13*, 429–435. [\[CrossRef\]](#)
83. Ke, H.M.; Lee, H.H.; Lin, C.I.; Liu, Y.C.; Lu, M.R.; Hsieh, J.A.; Chang, C.C.; Wu, P.H.; Lu, M.J.; Li, J.Y.; et al. *Mycena* genomes resolve the evolution of fungal bioluminescence. *Proc. Natl. Acad. Sci. USA* **2020**, *117*, 31267–31277. [\[CrossRef\]](#)
84. Ke, H.M.; Lu, M.R.; Chang, C.C.; Hsiao, C.; Ou, J.H.; Taneyama, Y.; Tsai, I.J. Evolution and Diversity of Bioluminescent Fungi. In *Evolution of Fungi and Fungal-Like Organisms*; Springer International Publishing: Cham, Switzerland, 2023; pp. 275–294.
85. Palkina, K.A.; Balakireva, A.V.; Belozerova, O.A.; Chepurnykh, T.V.; Markina, N.M.; Kovalchuk, S.I.; Tsarkova, A.S.; Mishin, A.S.; Yampolsky, I.V.; Sarkisyan, K.S. Domain truncation in hispidin synthase orthologs from non-bioluminescent fungi does not lead to hispidin biosynthesis. *Int. J. Mol. Sci.* **2023**, *24*, 1317. [\[CrossRef\]](#)
86. Kahlke, T.; Umbers, K.D.L. Bioluminescence. *Curr. Biol.* **2016**, *26*, R313–R314. [\[CrossRef\]](#)
87. Wilson, T.; Hastings, J.W. *Bioluminescence: Living Lights, Lights for Living*; Harvard University Press: Cambridge, MA, USA, 2013; p. 176. [\[CrossRef\]](#)
88. Liu, X.; Wang, M.; Liu, Y. Chemistry in fungal bioluminescence: Theoretical studies on biosynthesis of luciferin from caffeic acid and regeneration of caffeic acid from oxidized luciferin. *J. Fungi* **2023**, *9*, 369. [\[CrossRef\]](#) [\[PubMed\]](#)
89. Khakhar, A.; Starker, C.G.; Chamness, J.C.; Lee, N.; Stokke, S.; Wang, C.; Swanson, R.; Rizvi, F.; Imaizumi, T.; Voytas, D.F. Building customizable auto-luminescent luciferase-based reporters in plants. *eLife* **2020**, *9*, e52786. [\[CrossRef\]](#)
90. Dubois, R. Note sur la fonction photogénique chez la *Pholas dactylus*. *C. R. Seances Soc. Biol.* **1887**, *39*, e564.
91. Airth, R.L.; McElroy, W.D. Light emission from extract of luminous fungi. *J. Bacteriol.* **1959**, *77*, 249–250. Available online: <https://journals.asm.org/doi/pdf/10.1128/jb.77.2.249-250.1959> (accessed on 25 July 2024). [\[CrossRef\]](#)
92. Airth, R.L.; Foerster, G.E. The isolation of catalytic components required for cell-free fungal bioluminescence. *Arch. Biochem. Biophys.* **1962**, *97*, 567–573. [\[CrossRef\]](#) [\[PubMed\]](#)
93. Shimomura, O. *Bioluminescence: Chemical Principles and Methods*; World Scientific Publishing Co. Pte. Ltd.: Singapore, 2006; p. 500. [\[CrossRef\]](#)
94. Purtov, K.V.; Petushkov, V.N.; Baranov, M.S.; Mineev, K.S.; Rodionova, N.S.; Kaskova, Z.M.; Tsarkova, A.S.; Petunin, A.I.; Bondar, V.S.; Rodicheva, E.K.; et al. The chemical basis of fungal bioluminescence. *Angew. Chem. Int. Ed.* **2015**, *54*, 8124–8128. [\[CrossRef\]](#) [\[PubMed\]](#)
95. Wang, M.Y.; Liu, Y.J. Chemistry in fungal bioluminescence: A theoretical study from Luciferin to light emission. *J. Org. Chem.* **2021**, *86*, 1874–1881. [\[CrossRef\]](#) [\[PubMed\]](#)
96. Keller, N.P.; Turner, G.; Bennett, J.W. Fungal secondary metabolism—from biochemistry to genomics. *Nat. Rev. Microbiol.* **2005**, *3*, 937–947. [\[CrossRef\]](#) [\[PubMed\]](#)
97. Rokas, A.; Wisecaver, J.H.; Lind, A.L. The birth, evolution and death of metabolic gene clusters in fungi. *Nat. Rev. Microbiol.* **2018**, *16*, 731–744. [\[CrossRef\]](#)
98. Kim, J.; Park, M.J.; Shim, D.; Ryoo, R. De novo genome assembly of the bioluminescent mushroom *Omphalotus guepiniiformis* reveals an *Omphalotus*-specific lineage of the luciferase gene block. *Genomics* **2022**, *114*, e110514. [\[CrossRef\]](#) [\[PubMed\]](#)
99. Oliveira, A.G.; Stevani, C.V.; Waldenmaier, H.E.; Viviani, V.; Emerson, J.M.; Loros, J.J.; Dunlap, J.C. Circadian control sheds light on fungal bioluminescence. *Curr. Biol.* **2015**, *25*, 964–968. [\[CrossRef\]](#) [\[PubMed\]](#)
100. Mishara, M.; Srivastava, D. Bioluminescent fungi: Reviewing nature's riddle. *J. Mycopathol. Res.* **2021**, *59*, 199–206.
101. Sivinski, J.M. Arthropods attracted to luminous fungi. *Psyche A J. Entomol.* **1981**, *88*, 383–390. [\[CrossRef\]](#)
102. Bechara, E.J.H. Bioluminescence: A fungal nightlight with an internal timer. *Curr. Biol.* **2015**, *25*, R283–R285. [\[CrossRef\]](#) [\[PubMed\]](#)
103. Weinstein, P.; Delean, S.; Wood, T.; Austin, A.D. Bioluminescence in the ghost fungus *Omphalotus nidiformis* does not attract potential spore dispersing insects. *IMA Fungus* **2016**, *7*, 229–234. [\[CrossRef\]](#)
104. Lingle, W.L. Bioluminescence and ligninolysis during secondary metabolism in the fungus *Panellus stypticus*. *J. Biolum. Chemilum.* **1993**, *8*, e100.
105. Jabr, F. The Secret History of Bioluminescence. *Hakai Magazine*. 2016. Available online: <https://hakaimagazine.com/features/secret-history-bioluminescence/> (accessed on 14 July 2023).

106. Jain, N. In Meghalaya, Scientists Discover Glowing Mushrooms, Used by Locals as Natural Torches. 2020. Available online: <https://scroll.in/article/978927/in-meghalaya-scientists-discover-glowing-mushrooms-used-by-locals-as-natural-torches> (accessed on 14 July 2023).
107. Lepp, H. Aboriginal Use of Fungi. 2013. Available online: <https://www.anbg.gov.au/fungi/aboriginal.html> (accessed on 14 July 2023).
108. Araújo-Gomes, N.; Zambito, G.; Johnbosco, C.; Calejo, I.; Leijten, J.; Löwik, C.; Karperien, M.; Mezzanotte, L.; Teixeira, L.M. Bioluminescence imaging on-chip platforms for non-invasive high-content bioimaging. *Biosens. Bioelectron.* **2023**, *237*, e115510. [[CrossRef](#)] [[PubMed](#)]
109. Davies, K.A.; Welch, S.R.; Jain, S.; Sorvillo, T.E.; Coleman-McCray, J.D.; Montgomery, J.M.; Spiropoulou, C.F.; Albariño, C.; Spengler, J.R. Fluorescent and bioluminescent reporter mouse-adapted Ebola viruses maintain pathogenicity and can be visualized in vivo. *J. Infect. Dis.* **2023**, *228*, S536–S547. [[CrossRef](#)] [[PubMed](#)]
110. Li, B.; Chen, R.; Zhu, C.; Kong, F. Glowing plants can light up the night sky? A review. *Biotechnol. Bioeng.* **2021**, *118*, 3706–3715. [[CrossRef](#)]
111. Zheng, P.; Ge, J.; Ji, J.; Zhong, J.; Chen, H.; Luo, D.; Li, W.; Bi, B.; Ma, Y.; Tong, W.; et al. Metabolic engineering and mechanical investigation of enhanced plant autoluminescence. *Plant Biotechnol. J.* **2023**, *21*, 1671–1681. [[CrossRef](#)] [[PubMed](#)]
112. Steinberg, S.M.; Poziomek, E.J.; Engelmann, W.H.; Rogers, K.R. A review of environmental applications of bioluminescence measurements. *Chemosphere* **1995**, *30*, 2155–2197. [[CrossRef](#)]
113. Gianfreda, L.; Rao, M.A. Interactions between xenobiotics and microbial and enzymatic soil activity. *Crit. Rev. Environ. Sci. Technol.* **2008**, *3*, 269–310. [[CrossRef](#)]
114. Ventura, F.F.; Mendes, L.F.; Oliveira, A.G.; Bazito, R.C.; Bechara, E.J.; Freire, R.S.; Stevani, C.V. Evaluation of phenolic compound toxicity using a bioluminescent assay with the fungus *Gerronema viridilucens*. *Environ. Toxicol. Chem.* **2020**, *39*, 1558–1565. [[CrossRef](#)] [[PubMed](#)]
115. Ventura, F.F.; Soares, D.M.; Bayle, K.; Oliveira, A.G.; Bechara, E.J.; Freire, R.S.; Stevani, C.V. Toxicity of metal cations and phenolic compounds to the bioluminescent fungus *Neonothopanus gardneri*. *Environ. Adv.* **2021**, *4*, 100044. [[CrossRef](#)]
116. Endo, M.; Ozawa, T. Advanced bioluminescence system for in vivo imaging with brighter and red-shifted light emission. *Int. J. Mol. Sci.* **2020**, *21*, 6538. [[CrossRef](#)]
117. Chen, M.; Zhou, K.; Dai, S.Y.; Tadepalli, S.; Balakrishnan, P.B.; Xie, J.; Rami, F.E.; Dai, T.; Cui, L.; Idoyaga, J.; et al. In vivo bioluminescence imaging of granzyme B activity in tumor response to cancer immunotherapy. *Cell Chem. Biol.* **2022**, *29*, 1556–1567. [[CrossRef](#)] [[PubMed](#)]
118. Zhang, Q.; Song, B.; Xu, Y.; Yang, Y.; Ji, J.; Cao, W.; Lu, J.; Ding, J.; Cao, H.; Chu, B.; et al. In vivo bioluminescence imaging of natural bacteria within deep tissues via ATP-binding cassette sugar transporter. *Nat. Commun.* **2023**, *14*, 2331. [[CrossRef](#)] [[PubMed](#)]

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.